

長良サイエンス株式会社

Nagara Science Co., Ltd.

〒501-1121 岐阜市古市場 840

840 Furuichiba, Gifu 501-1121, JAPAN

Phone : +81-58-234-4257, Fax : +81-58-234-4724

E-mail : nagara@nsgifu.jp , http : //www.nsgifu.jp

Source	Code No.	Storage Conditions	Product Name · Purity · Molecular Formula = Molecular Weight · [CAS Registry Number]	Quantity · Price (JPY)
緑茶 Green tea	NH020102	2-10 °C	(-)-Epicatechin [(-)-EC] $\geq 99\%$ (HPLC)	10mg 8,000
	NH020103		$C_{15}H_{14}O_6 = 290.27$ [490-46-0]	100mg 44,000
	NH020202 NH020203	2-10 °C	(-)-Epigallocatechin [(-)-EGC] $\geq 99\%$ (HPLC)	10mg 12,000
			$C_{15}H_{14}O_7 = 306.27$ [970-74-1]	100mg 66,000
	NH020302 NH020303	2-10 °C	(-)-Epicatechin gallate [(-)-ECg] $\geq 99\%$ (HPLC)	10mg 12,000
			$C_{22}H_{18}O_{10} = 442.37$ [1257-08-5]	100mg 52,000
	NH020403	2-10 °C	(-)-Epigallocatechin gallate [(-)-EGCg] $\geq 98\%$ (HPLC)	100mg 12,000
			$C_{22}H_{18}O_{11} = 458.37$ [989-51-5]	
	NH020602 NH020603	2-10 °C	(-)-Epigallocatechin gallate [(-)-EGCg] $\geq 99\%$ (HPLC)	20mg 12,000
			$C_{22}H_{18}O_{11} = 458.37$ [989-51-5]	100mg 30,000
	NH020502 NH020503	2-10 °C	(+)-Catechin hydrate [(+)-C] $\geq 99\%$ (HPLC)	10mg 5,000
			$C_{15}H_{14}O_6 \cdot H_2O = 308.28$ [88191-48-4]	100mg 32,000
	NH021102	2-10 °C	(-)-Catechin [(-)-C] $\geq 98\%$ (HPLC)	10mg 23,000
			$C_{15}H_{14}O_6 = 290.27$ [18829-70-4]	
	NH021202	2-10 °C	(-)-Gallocatechin [(-)-GC] $\geq 98\%$ (HPLC)	10mg 34,000
			$C_{15}H_{14}O_7 = 306.27$ [3371-27-5]	
	NH021302	2-10 °C	(-)-Catechin gallate [(-)-Cg] $\geq 98\%$ (HPLC)	10mg 34,000
			$C_{22}H_{18}O_{10} = 442.37$ [130405-40-2]	
	NH021402	2-10 °C	(-)-Gallocatechin gallate [(-)-GCg] $\geq 98\%$ (HPLC)	10mg 23,000
			$C_{22}H_{18}O_{11} = 458.37$ [4233-96-9]	
	NH021502	2-10 °C	(+)-Epicatechin [(+)-EC] $\geq 98\%$ (HPLC)	10mg 22,000
			$C_{15}H_{14}O_6 = 290.27$ [35323-91-2]	
	NH022101	-20 °C	(-)-Epicatechin-3'-O-methylether [(-)-EC-3'-O-Me] (synthetic) $\geq 98\%$ (HPLC)	1mg 24,000
			$C_{16}H_{16}O_6 = 304.29$ [-]	
	NH022201	-20 °C	(-)-Epicatechin-4'-O-methylether [(-)-EC-4'-O-Me] (synthetic) $\geq 98\%$ (HPLC)	1mg 24,000
			$C_{16}H_{16}O_6 = 304.29$ [-]	
	NH022301	-20 °C	(-)-Epigallocatechin-3'-O-methylether [(-)-EGC-3'-O-Me] (synthetic) $\geq 98\%$ (HPLC)	1mg 24,000
			$C_{16}H_{16}O_7 = 320.29$ [-]	
	NH022401	-20 °C	(-)-Epigallocatechin-4'-O-methylether [(-)-EGC-4'-O-Me] (synthetic) $\geq 98\%$ (HPLC)	1mg 24,000
			$C_{16}H_{16}O_7 = 320.29$ [17291-05-3]	
	NH022501	-20 °C	(-)-Epicatechin-3'-O-methylether gallate [(-)-ECg-3'-O-Me] (synthetic) $\geq 98\%$ (HPLC)	1mg 36,000
			$C_{23}H_{20}O_{10} = 456.40$ [-]	
	NH022601	-20 °C	(-)-Epicatechin-4'-O-methylether gallate [(-)-ECg-4'-O-Me] (synthetic) $\geq 98\%$ (HPLC)	1mg 36,000
		$C_{23}H_{20}O_{10} = 456.40$ [-]		
NH022701	-20 °C	(-)-Epicatechin 3-(3'-O-methyl)gallate [(-)-ECg-3'-O-Me] (synthetic) $\geq 98\%$ (HPLC)	1mg 24,000	
		$C_{23}H_{20}O_{10} = 456.40$ [83104-86-3]		
NH027102	-20 °C	(-)-Epicatechin 3-(3'-O-methyl)gallate [(-)-ECg-3'-O-Me] (natural) $\geq 99\%$ (HPLC)	5mg 30,000	
		$C_{23}H_{20}O_{10} = 456.40$ [83104-86-3]		
NH022801	-20 °C	(-)-Epicatechin 3-(4'-O-methyl)gallate [(-)-ECg-4'-O-Me] (synthetic) $\geq 98\%$ (HPLC)	1mg 24,000	
		$C_{23}H_{20}O_{10} = 456.40$ [108907-44-4]		
NH022901	-20 °C	(-)-Epigallocatechin-3'-O-methylether gallate [(-)-EGCg-3'-O-Me] (synthetic) $\geq 98\%$ (HPLC)	1mg 36,000	
		$C_{23}H_{20}O_{11} = 472.40$ [298700-56-8]		
NH023001	-20 °C	(-)-Epigallocatechin-4'-O-methylether gallate [(-)-EGCg-4'-O-Me] (synthetic) $\geq 98\%$ (HPLC)	1mg 36,000	
		$C_{23}H_{20}O_{11} = 472.40$ [298700-57-9]		
NH027201 NH027202	-20 °C	(-)-Epigallocatechin 3-(3'-O-methyl)gallate [(-)-EGCg-3'-O-Me] (natural) $\geq 99\%$ (HPLC)	2mg 20,000	
		$C_{23}H_{20}O_{11} = 472.40$ [83104-87-4]	10mg 30,000	
NH023201	-20 °C	(-)-Epigallocatechin 3-(4'-O-methyl)gallate [(-)-EGCg-4'-O-Me] (synthetic) $\geq 98\%$ (HPLC)	1mg 24,000	
		$C_{23}H_{20}O_{11} = 472.40$ [224434-07-5]		
NH023301	-20 °C	(-)-Gallocatechin 3-(3'-O-methyl)gallate [(-)-GCg-3'-O-Me] (synthetic) $\geq 98\%$ (HPLC)	1mg 100,000	
		$C_{23}H_{20}O_{11} = 472.40$ [264147-80-0]		
NH023401	-20 °C	(-)-Gallocatechin 3-(4'-O-methyl)gallate [(-)-GCg-4'-O-Me] (synthetic) $\geq 99\%$ (HPLC)	1mg 100,000	
		$C_{23}H_{20}O_{11} = 472.40$ [-]		
NH023501	-20 °C	(-)-EGCg-3', 4'-di-O-Me $\geq 99\%$ (HPLC)	1mg 50,000	
		$C_{24}H_{22}O_{11} = 486.42$ [-]		
NH023601	-20 °C	(-)-EGCg-4', 4'-di-O-Me $\geq 99\%$ (HPLC)	1mg 50,000	
		$C_{24}H_{22}O_{11} = 486.42$ [-]		
次ページに続く	NH023701	-20 °C	(-)-EGCg-3', 4'-di-O-Me $\geq 99\%$ (HPLC)	1mg 50,000
			$C_{24}H_{22}O_{11} = 486.42$ [-]	

Source	Code No.	Storage Conditions	Product Name · Purity · Molecular Formula=Molecular Weight · [CAS Registry Number]	Quantity · Price (JPY)
緑茶 Green tea	NH025104	2-10 °C	Catechin Mixture { (-)-Epicatechin, (-)-Epigallocatechin, (-)-Epicatechin gallate, (-)-Epigallocatechin gallate }, from Green tea ≥Total 80% (HPLC)	1g 19,800
	NH026102	2-10 °C	Strictinin C ₂₇ H ₂₂ O ₁₈ = 634.45 [517-46-4] ≥98% (HPLC)	5mg 30,000
	NS630102 <i>new</i>	2-10 °C	Quercetin 3- <i>O</i> -β-D-glucosyl-(1→3)-L-rhamnosyl-(1→6)-β-D-glucoside C ₃₃ H ₄₀ O ₂₁ = 772.66 [136449-10-0] ≥98% (HPLC)	5mg 40,000
	NS630202 <i>new</i>	2-10 °C	Quercetin 3- <i>O</i> -β-D-glucosyl-(1→3)-α-L-rhamnosyl-(1→6)-β-D-galactoside C ₃₃ H ₄₀ O ₂₁ = 772.66 [134953-93-8] ≥98% (HPLC)	5mg 60,000
茶サポゲニン	NS590102 <i>new</i>	2-10 °C	R ₁ -Barrigenol C ₃₀ H ₅₀ O ₆ = 506.72 [15399-43-6] ≥99% (HPLC)	5mg 36,000
紅茶 Black tea	NH030101	-20 °C	Theaflavin C ₂₀ H ₂₁ O ₁₂ = 564.49 [4670-05-7] ≥98% (HPLC)	1mg 12,000
	NH030102	-20 °C	Theaflavin C ₂₀ H ₂₁ O ₁₂ = 564.49 [4670-05-7] ≥98% (HPLC)	5mg (新容量) 24,000
	NH030201	-20 °C	Theaflavin 3- <i>O</i> -gallate C ₃₀ H ₂₈ O ₁₆ = 716.60 [30462-34-1] ≥98% (HPLC)	1mg 15,000
	NH030202	-20 °C	Theaflavin 3- <i>O</i> -gallate C ₃₀ H ₂₈ O ₁₆ = 716.60 [30462-34-1] ≥98% (HPLC)	5mg (新容量) 30,000
	NH030301	-20 °C	Theaflavin 3'- <i>O</i> -gallate C ₃₀ H ₂₈ O ₁₆ = 716.60 [28543-07-9] ≥98% (HPLC)	1mg 15,000
NH030302	-20 °C	Theaflavin 3'- <i>O</i> -gallate C ₃₀ H ₂₈ O ₁₆ = 716.60 [28543-07-9] ≥98% (HPLC)	5mg (新容量) 30,000	
NH030401	-20 °C	Theaflavin 3,3'-di- <i>O</i> -gallate C ₄₃ H ₃₂ O ₂₀ = 868.70 [30462-35-2] ≥98% (HPLC)	1mg 15,000	
NH030402	-20 °C	Theaflavin 3,3'-di- <i>O</i> -gallate C ₄₃ H ₃₂ O ₂₀ = 868.70 [30462-35-2] ≥98% (HPLC)	5mg (新容量) 30,000	
大豆 Soy bean	NH010102	冷暗所	Daidzein C ₁₅ H ₁₀ O ₄ = 254.24 [486-66-8] ≥98% (HPLC)	10mg 5,000
	NH010103	冷暗所	Daidzein C ₁₅ H ₁₀ O ₄ = 254.24 [486-66-8] ≥98% (HPLC)	100mg 29,800
	NH010202	冷暗所	Glycitein C ₁₆ H ₁₂ O ₅ = 284.26 [40957-83-3] ≥98% (HPLC)	10mg 13,500
	NH010203	冷暗所	Glycitein C ₁₆ H ₁₂ O ₅ = 284.26 [40957-83-3] ≥98% (HPLC)	100mg 98,000
	NH010302	冷暗所	Genistein C ₁₅ H ₁₀ O ₅ = 270.24 [446-72-0] ≥98% (HPLC)	10mg 5,000
	NH010303	冷暗所	Genistein C ₁₅ H ₁₀ O ₅ = 270.24 [446-72-0] ≥98% (HPLC)	100mg 19,000
	NH012102	冷暗所	Daidzin C ₂₁ H ₂₀ O ₉ = 416.38 [552-66-9] ≥99% (HPLC) 純度規格変更	10mg 12,000
	NH012103	冷暗所	Daidzin C ₂₁ H ₂₀ O ₉ = 416.38 [552-66-9] ≥99% (HPLC) 純度規格変更	100mg 98,000
	NH012202	冷暗所	Glycitin C ₂₂ H ₂₂ O ₁₀ = 446.40 [40246-10-4] ≥99% (HPLC) 純度規格変更	10mg 13,500
	NH012203	冷暗所	Glycitin C ₂₂ H ₂₂ O ₁₀ = 446.40 [40246-10-4] ≥99% (HPLC) 純度規格変更	100mg 98,000
	NH012302	冷暗所	Genistin C ₂₁ H ₂₀ O ₁₀ = 432.38 [529-59-9] ≥99% (HPLC) 純度規格変更	10mg 13,500
	NH012303	冷暗所	Genistin C ₂₁ H ₂₀ O ₁₀ = 432.38 [529-59-9] ≥99% (HPLC) 純度規格変更	100mg 98,000
	NH013101	-20 °C	6'- <i>O</i> -Acetyldaidzin C ₂₃ H ₂₂ O ₁₀ = 458.41 [71385-83-6] ≥98% (HPLC)	1mg 15,000
	NH013201	-20 °C	6'- <i>O</i> -Acetylglycitin C ₂₄ H ₂₄ O ₁₁ = 488.44 [134859-96-4] ≥98% (HPLC)	1mg 15,000
	NH013301	-20 °C	6'- <i>O</i> -Acetylgenistin C ₂₃ H ₂₂ O ₁₁ = 474.41 [73566-30-0] ≥98% (HPLC)	1mg 15,000
	NH014101	-20 °C	6'- <i>O</i> -Malonyldaidzin C ₂₄ H ₂₂ O ₁₂ = 502.42 [124590-31-4] ≥98% (HPLC)	1mg 15,000
	NH014201	-20 °C	6'- <i>O</i> -Malonylglycitin C ₂₅ H ₂₄ O ₁₃ = 532.45 [137705-39-6] ≥98% (HPLC)	1mg 15,000
	NH014301	-20 °C	6'- <i>O</i> -Malonylgenistin C ₂₄ H ₂₂ O ₁₃ = 518.42 [51011-05-3] ≥98% (HPLC)	1mg 15,000
NH015204	冷暗所	Isoflavone Aglycon Mixture B (Genistein ≥ 50%), from Soybean ≥Total 95% (HPLC)	1g 29,800	
NH016104	冷暗所	Isoflavone Glucoside Mixture A (Daidzin ≥ 50%), from Soybean ≥Total 95% (HPLC)	1g 30,000	
NH017102	2-10 °C	8-Hydroxydaidzein C ₁₅ H ₁₀ O ₅ = 270.24 [75187-63-2] ≥98% (HPLC)	5mg 36,000	
NH017202	2-10 °C	8-Hydroxyglycitein C ₁₆ H ₁₂ O ₆ = 300.26 [113762-90-6] ≥98% (HPLC)	5mg 36,000	
NH017302	2-10 °C	8-Hydroxygenistein C ₁₅ H ₁₀ O ₆ = 286.24 [13539-27-0] ≥98% (HPLC)	5mg 36,000	

Source	Code No.	Storage Conditions	Product Name · Purity · Molecular Formula = Molecular Weight · [CAS Registry Number]	Quantity · Price (JPY)
ウコン Turmeric	NH040103	2-10 °C	Curcumin 1 (Curcumin) $\geq 98\%$ (HPLC) C ₂₁ H ₂₀ O ₆ = 368.38 [458-37-7, 147556-16-9]	100mg 15,000
	NH040202	2-10 °C	Curcumin 2 $\geq 98\%$ (HPLC) C ₂₀ H ₁₈ O ₅ = 338.35 [22608-11-3, 300853-82-1, 478919-22-1]	10mg 12,000
	NH040302	2-10 °C	Curcumin 3 $\geq 98\%$ (HPLC) C ₁₉ H ₁₆ O ₄ = 308.33 [33171-05-0, 52328-96-8]	10mg 12,000
	NH040402	2-10 °C	Cyclocurcumin (synthetic) $\geq 99\%$ (HPLC) C ₂₁ H ₂₀ O ₆ = 368.38 [153127-42-5]	5mg 30,000
	NH040702	2-10 °C	Dihydrocurcumin $\geq 99\%$ (HPLC) C ₂₁ H ₂₂ O ₆ = 370.40 [76474-56-1, 943330-36-7]	5mg 24,000
	NH040802	2-10 °C	Tetrahydrocurcumin (synthetic) $\geq 99\%$ (HPLC) C ₂₁ H ₂₁ O ₆ = 372.41 [36062-04-1, 426828-40-2]	5mg 12,000
	NH040502	2-10 °C	Tetrahydrodemethoxycurcumin (synthetic) $\geq 99\%$ (HPLC) C ₂₀ H ₂₂ O ₅ = 342.39 [149579-07-7, 426828-41-3]	5mg 24,000
	NH040602	2-10 °C	Tetrahydrobisdemethoxycurcumin (synthetic) $\geq 99\%$ (HPLC) C ₁₉ H ₂₁ O ₄ = 312.36 [113482-94-3, 426828-42-4]	5mg 24,000
	NH042101 NH042102	-20 °C	Bisacurone $\geq 99\%$ (HPLC) C ₁₅ H ₂₁ O ₃ = 252.35 [120681-81-4]	2mg 32,000 5mg 60,000
	NH0422	-20 °C	Bisacurone A $\geq 99\%$ (HPLC) C ₁₅ H ₂₁ O ₃ = 252.35 [127214-84-0]	要問合せ
	NH0423	-20 °C	Bisacurone B $\geq 99\%$ (HPLC) C ₁₅ H ₂₁ O ₃ = 252.35 [127214-85-1]	要問合せ
	NH0424	-20 °C	1-Bisabolone $\geq 99\%$ (HPLC) C ₁₅ H ₂₁ O = 220.35 [61432-71-1]	要問合せ
	NH0425	-20 °C	Bisacurool $\geq 99\%$ (HPLC) C ₁₅ H ₂₁ O = 220.35 [120681-80-3]	要問合せ
	NH042602	-20 °C	Curlone $\geq 95\%$ (HPLC) C ₁₅ H ₂₂ O = 218.33 [87440-60-6]	5mg 36,000
	NH0427	-20 °C	2-Hepten-4-one, 5-hydroxy-2-methyl-6-(4-methylphenyl)- $\geq 99\%$ (HPLC) C ₁₅ H ₂₀ O ₂ = 232.32 [192131-18-3]	要問合せ
	NH042802	-20 °C	<i>ar</i> -Turmerone $\geq 99\%$ (HPLC) C ₁₅ H ₂₀ O = 216.32 [532-65-0]	5mg 36,000
	NH042901	-20 °C	α -Turmerone $\geq 95\%$ (HPLC) C ₁₅ H ₂₂ O = 218.33 [82508-15-4]	1mg 36,000
	NH043001	-20 °C	Turmeronol A $\geq 99\%$ (HPLC) C ₁₅ H ₂₁ O ₂ = 232.32 [131651-37-1]	2mg 50,000
	NH043101	-20 °C	Turmeronol B $\geq 99\%$ (HPLC) C ₁₅ H ₂₁ O ₂ = 232.32 [131651-38-2]	2mg 60,000
	イチョウ Ginkgo biloba	NH050102 NH050103	2-10 °C	Ginkgolide A $\geq 98\%$ (HPLC) C ₂₀ H ₂₁ O ₉ = 408.40 [15291-75-5]
NH050202 NH050203		2-10 °C	Ginkgolide B $\geq 98\%$ (HPLC) C ₂₀ H ₂₁ O ₁₀ = 424.40 [15291-77-7]	10mg 8,000 100mg 56,000
NH050302 NH050303		2-10 °C	Ginkgolide C $\geq 98\%$ (HPLC) C ₂₀ H ₂₁ O ₁₁ = 440.40 [15291-76-6]	10mg 8,000 100mg 56,000
NH050402 NH050403		2-10 °C	(-)-Bilobalide $\geq 98\%$ (HPLC) C ₁₅ H ₁₈ O ₈ = 326.30 [33570-04-6]	10mg 12,000 100mg 84,000
NH052102		2-10 °C	Ginkgolic acid 15:1 [Anacardic acid 15:1, Ginkgolic acid I] $\geq 99\%$ (HPLC) C ₂₂ H ₃₁ O ₃ = 346.50 [22910-60-7]	10mg 29,800
NH052202		2-10 °C	Ginkgolic acid 17:1 [Anacardic acid 17:1, Ginkgolic acid II] $\geq 99\%$ (HPLC) C ₂₄ H ₃₃ O ₃ = 374.56 [111047-30-4]	10mg 29,800
NH052302		2-10 °C	Ginkgolic acid 13:0 [Anacardic acid 13:0] $\geq 99\%$ (HPLC) C ₂₀ H ₂₉ O ₃ = 320.47 [20261-38-5]	5mg 29,800
NH052402 NH052403		2-10 °C	Ginkgolic acid 15:0 [Anacardic acid 15:0] $\geq 99\%$ (HPLC) C ₂₂ H ₃₁ O ₃ = 348.52 [16611-84-0]	10mg 18,000 100mg 126,000
NH053102		2-10 °C	Ginkgolic acid 15:1+Ginkgolic acid 13:0 (4:1) [Anacardic acid 15:1+Anacardic acid 13:0] $\geq 99\%$ (HPLC)	10mg 24,000
NH053202		2-10 °C	Ginkgolic acid 17:1+Ginkgolic acid 15:0 (6:1) [Anacardic acid 17:1+Anacardic acid 15:0] $\geq 99\%$ (HPLC)	10mg 24,000
NH053303 NH053304		2-10 °C	Ginkgolic acids [Anacardic acids] { Ginkgolic acid 13:0, Ginkgolic acid 15:0, Ginkgolic acid 15:1, Ginkgolic acid 17:1 etc. }, from Ginkgo biloba \geq Total 95% (HPLC)	100mg 15,000 1g 30,000

Source	Code No.	Storage Conditions	Product Name · Purity · Molecular Formula = Molecular Weight · [CAS Registry Number]	Quantity · Price (JPY)
紫根 Lithospermum erythrorhizon	NH060002	2-10 °C	Shikonin $\geq 99\%$ ee (HPLC) C ₁₆ H ₁₆ O ₅ = 288.30 [517-89-5]	10mg 30,000
	NH060102 NH060103 NH060104	2-10 °C	Shikonin (Shikonin/Alkannin ≒ 6 : 1) \geq Total 98% (HPLC) C ₁₆ H ₁₆ O ₅ = 288.30 [517-89-5, 517-88-4]	10mg 14,000 100mg 32,000 1g 200,000
	NH063202	2-10 °C	Shikalkin (Shikonin/Alkannin ≒ 1 : 1) \geq Total 98% (HPLC) C ₁₆ H ₁₆ O ₅ = 288.30 [54952-43-1]	10mg 22,000
	NH063102	2-10 °C	Alkannin $\geq 99\%$ ee (HPLC) C ₁₆ H ₁₆ O ₅ = 288.30 [517-88-4]	10mg 30,000
	NH060202	2-10 °C	Acetylshikonin (natural) $\geq 98\%$ (HPLC) C ₁₈ H ₁₈ O ₆ = 330.33 [24502-78-1, 34232-27-4]	10mg 24,000
	NH060302	2-10 °C	Isobutyrylshikonin (natural) $\geq 98\%$ (HPLC) C ₂₀ H ₂₀ O ₆ = 358.39 [52438-12-7, 87562-78-5]	10mg 24,000
	NH060402	2-10 °C	β -Hydroxyisovalerylshikonin (natural) $\geq 98\%$ (HPLC) C ₂₁ H ₂₁ O ₇ = 388.41 [7415-78-3, 87798-74-1]	10mg 24,000
	NH060501	2-10 °C	Isovalerylshikonin (natural) $\geq 98\%$ (HPLC) C ₂₁ H ₂₁ O ₆ = 372.41 [52387-14-1, 69175-71-9]	1mg 24,000
	NH060601	2-10 °C	α -Methyl- <i>n</i> -butyrylshikonin (natural) $\geq 98\%$ (HPLC) C ₂₁ H ₂₁ O ₆ = 372.41 [92175-42-3, 92175-40-1]	1mg 24,000
	NH060702	2-10 °C	β , β -Dimethylacrylshikonin (β , β -Dimethylacrylalkannin) (natural) $\geq 98\%$ (HPLC) C ₂₁ H ₂₂ O ₆ = 370.40 [34539-65-6]	10mg 28,000
	NH060901	2-10 °C	β -Acetoxyisovalerylshikonin (β -Acetoxyisovalerylalkannin) (natural) $\geq 98\%$ (HPLC) C ₂₄ H ₂₄ O ₈ = 430.45 [98936-77-7, 69091-17-4]	2mg 36,000
ローヤルゼリー Royal jelly	NH070102 NH070103	2-10 °C	(<i>E</i>)-10-Hydroxy-2-decenoic acid $\geq 99\%$ (HPLC) C ₁₀ H ₁₈ O ₃ = 186.25 [14113-05-4] *第十八改正日本薬局方第二追補 一般試験法 試薬・試液 10-ヒドロキシ-2-(<i>E</i>)-デセン酸, 定量用及び薄層クロマトグラフィー用に適合しています。 *弊社追加規格: $\geq 99\%$ (HPLC)	10mg 8,000 100mg 64,000
甘草 Licorice (Glycyrrhiza)	NH080102 NH080103	2-10 °C	Glycyrrhizin (Glycyrrhizic acid) $\geq 99\%$ (HPLC) C ₄₂ H ₆₂ O ₁₆ · <i>n</i> H ₂ O = 822.93 + [103000-77-7, 1405-86-3]	20mg 7,000 100mg 22,000
	NH080202 NH080203	冷暗所	Glycyrrhizic acid monoammonium salt <i>n</i> -hydrate $\geq 99\%$ (HPLC) C ₄₂ H ₆₁ O ₁₆ NH ₄ · <i>n</i> H ₂ O = 839.96 + [691358-65-3, 53956-04-0]	10mg 12,000 100mg 84,000
	NH080302 NH080303	冷暗所	Glycyrrhizic acid dipotassium salt $\geq 99\%$ (HPLC) C ₄₂ H ₆₀ K ₂ O ₁₆ = 899.11 [1007893-70-0, 68797-35-3]	10mg 12,000 100mg 84,000
	NH080502	冷暗所	Glycyrrhetic acid 3- <i>O</i> -glucuronide $\geq 99\%$ (HPLC) C ₃₆ H ₅₄ O ₁₀ = 646.82 [34096-83-8]	10mg 28,000
	NH080402 NH080403	冷暗所	18 β -Glycyrrhetic acid (18 β -Glycyrrhetic acid) $\geq 99\%$ (HPLC) C ₃₀ H ₄₆ O ₄ = 470.68 [471-53-4]	10mg 7,000 100mg 49,000
	NH080602	2-10 °C	18 α -Glycyrrhetic acid (18 α -Glycyrrhetic acid) $\geq 98\%$ (HPLC) C ₃₀ H ₄₆ O ₄ = 470.68 [1449-05-4]	10mg 20,000
	NH080702	2-10 °C	Licoricesaponin G ₂ (24-Hydroxyglycyrrhizin) $\geq 98\%$ (HPLC) C ₄₂ H ₆₂ O ₁₇ = 838.93 [118441-84-2]	10mg 20,000
	NH080802	2-10 °C	24-Hydroxyglycyrrhetic acid $\geq 98\%$ (HPLC) C ₃₀ H ₄₆ O ₅ = 486.68 [52911-55-4]	5mg 45,000
	NH080902	2-10 °C	18 α -24-Hydroxyglycyrrhetic acid $\geq 98\%$ (HPLC) C ₃₀ H ₄₆ O ₅ = 486.68 [-]	5mg 67,500
	NH081002	2-10 °C	Licoricesaponin H ₂ $\geq 98\%$ (HPLC) C ₄₂ H ₆₂ O ₁₆ = 822.93 [118441-85-3]	10mg 30,000
	NH081102	2-10 °C	Liquiritic acid $\geq 98\%$ (HPLC) C ₃₀ H ₄₆ O ₄ = 470.68 [10379-72-3]	5mg 45,000
	NH081202	2-10 °C	18 α -Liquiritic acid $\geq 98\%$ (HPLC) C ₃₀ H ₄₆ O ₄ = 470.68 [10379-78-9]	5mg 67,500

Source	Code No.	Storage Conditions	Product Name · Purity · Molecular Formula = Molecular Weight · [CAS Registry Number]	Quantity · Price (JPY)
五味子 Schisandra fruit	NH090102	2-10 °C	Schizandrin C ₂₁ H ₃₂ O ₇ = 432.51 [7432-28-2] ≥99% (HPLC)	10mg 29,800
	NH090202	2-10 °C	Gomisin A C ₂₃ H ₂₈ O ₇ = 416.46 [58546-54-6] ≥99% (HPLC)	10mg 26,000
	NH090302	2-10 °C	Gomisin N C ₂₃ H ₂₈ O ₆ = 400.46 [69176-52-9] ≥99% (HPLC)	10mg 29,800
	NH090402	2-10 °C	Deoxyschizandrin C ₂₁ H ₃₂ O ₆ = 416.51 [61281-38-7] ≥99% (HPLC)	5mg 29,800
	NH090502	2-10 °C	(±)-γ-Schizandrin (Schizandrin B) C ₂₃ H ₂₈ O ₆ = 400.46 [61281-37-6] ≥99% (HPLC)	5mg 29,800
	NH090602	2-10 °C	Wuweizisu C C ₂₂ H ₂₄ O ₆ = 384.42 [61301-33-5] ≥99% (HPLC)	5mg 29,800
唐辛子 Red pepper	NH100102	2-10 °C	Capsaicin ≥99% (HPLC phenylcolumn) C ₁₈ H ₂₇ NO ₃ = 305.41 [404-86-4]	10mg 17,000
	NH100103			100mg 119,000
	NH102102	2-10 °C	Capsaicin ≥95% (HPLC)	50mg 20,000
	NH102103			500mg 140,000
	NH100202	2-10 °C	Dihydrocapsaicin ≥99% (HPLC phenylcolumn) C ₁₈ H ₂₉ NO ₃ = 307.43 [19408-84-5]	10mg 20,000
	NH100203			100mg 140,000
	NH100301	2-10 °C	Nordihydrocapsaicin ≥95% (HPLC) C ₁₇ H ₂₇ NO ₃ = 293.40 [28789-35-7]	1mg 28,000
NH100401	2-10 °C	N-Vanillylnonanamide ≥95% (HPLC) C ₁₇ H ₂₇ NO ₃ = 293.40 [2444-46-4]	1.0mg 28,000	
NH100501	2-10 °C	N-Vanillyldecanamide ≥95% (HPLC) C ₁₈ H ₂₉ NO ₃ = 307.43 [31078-36-1]	1.0mg 28,000	
NH100602	2-10 °C	N-Vanillylnonanamide [Capsaicin (synthetic)] ≥99% (HPLC)	20mg 20,000	
NH100603			100mg 30,000	
丁子 Clove	NS120102	2-10 °C	Clove 3 C ₁₆ H ₁₈ O ₉ = 354.31 [152041-16-2] ≥95% (HPLC)	5mg 30,000
	NS120202	2-10 °C	Biflorin C ₁₆ H ₁₈ O ₉ = 354.31 [89701-85-9] ≥95% (HPLC)	5mg 30,000
	NS122102	2-10 °C	Casuarictin C ₄₁ H ₂₈ O ₂₆ = 936.65 [79786-00-8] ≥95% (HPLC)	5mg 30,000
	NS122202	2-10 °C	Eugenin C ₄₁ H ₃₀ O ₂₆ = 938.66 [81571-72-4] ≥95% (HPLC)	5mg 30,000
	NS122302	2-10 °C	Tellimagrandin I C ₃₄ H ₂₆ O ₂₂ = 786.56 [79786-08-6 · 118014-28-1 · 118014-29-2] ≥95% (HPLC)	5mg 30,000
	NS122402	2-10 °C	1,3-Di-O-galloyl-4,6-O-(S)-hexahydroxydiphenyl-β-D-glucose ≥90% (HPLC) C ₃₄ H ₂₆ O ₂₂ = 786.56 [139975-10-3 · 1630724-71-8]	1mg 30,000
チョウトウコウ Uncaria rhynchophylla	NS130102	2-10 °C	Rhynchophylline C ₂₂ H ₂₈ N ₂ O ₄ = 384.47 [76-66-4] ≥99% (HPLC)	10mg 39,800
除虫菊	NS140102	-20 °C	Pyrethrin I C ₂₁ H ₂₈ O ₃ = 328.45 [121-21-1] ≥95% (HPLC)	10mg 64,000
	NS140202	-20 °C	Pyrethrin II C ₂₂ H ₂₈ O ₅ = 372.45 [121-29-9] ≥95% (HPLC)	10mg 64,000
	NS140302	-20 °C	Cinerin I C ₂₀ H ₂₈ O ₃ = 316.43 [25402-06-6] ≥95% (HPLC)	10mg 64,000
	NS140402	-20 °C	Cinerin II C ₂₁ H ₂₈ O ₅ = 360.44 [121-20-0] ≥95% (HPLC)	10mg 64,000
	NS140502	-20 °C	Jasmolin I C ₂₁ H ₃₀ O ₃ = 330.46 [4466-14-2] ≥95% (HPLC)	5mg 64,000
	NS140602	-20 °C	Jasmolin II C ₂₂ H ₃₀ O ₅ = 374.47 [1172-63-0] ≥95% (HPLC)	5mg 64,000
カンキツ ヘスペリジン類	NS151102	2-10 °C	(2S)-Hesperidin C ₂₈ H ₃₄ O ₁₅ = 610.56 [520-26-3] ≥99% (HPLC) (98% ee)	10mg 50,000
	NS151202	2-10 °C	(2R)-Hesperidin C ₂₈ H ₃₄ O ₁₅ = 610.56 [369593-42-0] ≥99% (HPLC) (98% ee)	10mg 50,000
	NS153102	2-10 °C	(2S)-Naringin C ₂₇ H ₃₂ O ₁₄ = 580.53 [10236-47-2] ≥99% (HPLC) (98% ee)	10mg 50,000
	NS153202	2-10 °C	(2R)-Naringin C ₂₇ H ₃₂ O ₁₄ = 580.53 [58001-41-5] ≥99% (HPLC) (95% ee)	10mg 50,000

Source et. al.	Code No.	Storage Conditions	Product Name · Purity · Molecular Formula = Molecular Weight · [CAS Registry Number]	Quantity · Price (JPY)	
ステビア Stevia	NS160303	2-10 °C	Stevioside C ₃₈ H ₆₀ O ₁₈ =804.87 [57817-89-7] ≥99% (HPLC)	100mg 15,000	
	NS160402	2-10 °C	Dulcoside A C ₃₈ H ₆₀ O ₁₇ =788.87 [64432-06-0] ≥98% (HPLC)	2mg 30,000	
	NS160503	2-10 °C	Rebaudioside A C ₄₄ H ₇₀ O ₂₃ =967.01 [58543-16-1] ≥99% (HPLC)	100mg 15,000	
	NS160702	2-10 °C	Rebaudioside C C ₄₄ H ₇₀ O ₂₂ =951.01 [63550-99-2] ≥98% (HPLC)	5mg 36,000	
胡麻 Sesamum indicum	NS180103 NS180104	-20 °C	(+)-Sesamin C ₂₀ H ₁₈ O ₆ = 354.35 [607-80-7] ≥99% (HPLC)	100mg 20,000 1g 60,000	
	NS181102 NS181103	-20 °C	(+)-Sesamolin C ₂₀ H ₁₈ O ₇ = 370.35 [526-07-8] ≥99% (HPLC)	10mg 20,000 100mg 100,000	
	NS180202 NS180203	-20 °C	(+)-Episesamin C ₂₀ H ₁₈ O ₆ = 354.35 [133-03-9] ≥99% (HPLC)	10mg 30,000 100mg 210,000	
	NS180301	-20 °C	Diasesamin C ₂₀ H ₁₈ O ₆ = 354.35 [551-30-4] ≥99% (HPLC)	2mg 50,000	
	NS182102	-20 °C	Sesaminol C ₂₀ H ₁₈ O ₇ = 370.35 [74061-79-3] ≥99% (HPLC)	10mg 36,000	
	NS182202	-20 °C	2-Episesaminol C ₂₀ H ₁₈ O ₇ = 370.35 [104319-96-2] ≥98% (HPLC)	10mg 36,000	
	NS182301	-20 °C	6-Episesaminol C ₂₀ H ₁₈ O ₇ = 370.35 [105616-55-5] ≥99% (HPLC)	2mg 36,000	
	NS182401	-20 °C	Diasesaminol C ₂₀ H ₁₈ O ₇ = 370.35 [110300-35-1] ≥98% (HPLC)	2mg 70,000	
	NS183102	-20 °C	Samin C ₁₃ H ₁₄ O ₅ = 250.25 [166239-82-3] ≥99% (HPLC)	10mg 36,000	
	NS183202	-20 °C	Disaminyl ether C ₂₃ H ₂₃ O ₉ = 482.48 [30186-93-7] ≥99% (HPLC)	5mg 36,000	
	NS184103 NS184104	-20 °C	Sesamol C ₇ H ₆ O ₃ = 138.12 [533-31-3] ≥99% (HPLC)	100mg 12,000 1g 20,000	
	NS185102	-20 °C	Sesaminol triglucoside C ₃₈ H ₄₈ O ₂₂ = 856.77 [157469-83-5] ≥99% (HPLC)	5mg 36,000	
	NS185201	-20 °C	Sesaminol (1→2) diglucoside C ₃₂ H ₃₈ O ₁₇ = 694.63 [157469-82-4] ≥99% (HPLC)	1mg 36,000	
	NS185301	-20 °C	Sesaminol (1→6) diglucoside C ₃₂ H ₃₈ O ₁₇ = 694.63 [474431-66-8] ≥99% (HPLC)	1mg 36,000	
	NS185401 <i>new</i>	-20 °C	Sesaminol monoglucoside C ₂₆ H ₂₈ O ₁₂ = 532.49 [153512-13-1] ≥99% (HPLC)	1mg 70,000	
	NS1855 <i>new</i>	-20 °C	Sesamolignol diglucoside C ₃₂ H ₄₀ O ₁₇ = 696.65 [749874-95-1] ≥99% (HPLC)	お問い合わせください	
	NS1856 <i>new</i>	-20 °C	Sesamolignol monoglucoside C ₂₆ H ₃₀ O ₁₂ = 534.51 [1353018-00-4] ≥99% (HPLC)	お問い合わせください	
	細辛 Asiasarum sieboldii	NS190102	-20 °C	(-)-Asarinin C ₂₀ H ₁₈ O ₆ = 354.35 [133-04-0] ≥99% (HPLC)	10mg 44,000
		NS190202	-20 °C	(-)-Sesamin/(+)-Sesamin [natural enantiomeric mixture] ≥Total 99% (HPLC) C ₂₀ H ₁₈ O ₆ = 354.35	5mg 44,000
NS190302		-20 °C	(-)-Sesamin C ₂₀ H ₁₈ O ₆ = 354.35 [13079-95-3] ≥99% (HPLC)	2mg 44,000	
NS191102		-20 °C	(±)-Car-3-ene-2,5-dione C ₁₀ H ₁₂ O ₂ = 164.20 [6617-34-1] ≥99% (HPLC)	5mg 44,000	
亜麻仁 Flaxseed	NS200002 NS200003	-20 °C	SDG Mixture C ₃₂ H ₄₆ O ₁₆ = 686.70 ≥Total 99% (HPLC)	10mg 40,000 100mg 320,000	
	NS200102 NS200103	-20 °C	SDG Mixture C ₃₂ H ₄₆ O ₁₆ = 686.70 ≥Total 95% (TLC)	10mg 20,000 100mg 160,000	
	NS200202	-20 °C	(2 <i>R</i> ,3 <i>R</i>)-SDG C ₃₂ H ₄₆ O ₁₆ = 686.70 [158932-33-3] ≥99% (HPLC)	5mg 40,000	
	NS200302	-20 °C	(2 <i>R</i> ,3 <i>S</i>)-SDG C ₃₂ H ₄₆ O ₁₆ = 686.70 [492449-92-0] ≥99% (HPLC)	2mg 40,000	

Source et. al.	Code No.	Storage Conditions	Product Name · Purity · Molecular Formula = Molecular Weight · [CAS Registry Number]	Quantity · Price (JPY)
ブクリョウ	NS230102	-20 °C	Pachymic acid C ₃₃ H ₅₂ O ₅ = 528.76 [29070-92-6] ≥99% (HPLC)	10mg 36,000
	NS230201	-20 °C	Eburicoic acid C ₃₁ H ₅₀ O ₃ = 470.73 [560-66-7] ≥99% (HPLC)	2mg 36,000
	NS230302	-20 °C	Dehydroeburicoic acid C ₃₁ H ₄₈ O ₃ = 468.71 [6879-05-6] ≥99% (HPLC)	5mg 36,000
	NS230402	-20 °C	Dehydropachymic acid C ₃₃ H ₅₀ O ₅ = 526.75 [77012-31-8] ≥99% (HPLC)	5mg 50,000
	NS230502	-20 °C	Trametenolic acid C ₃₀ H ₄₈ O ₃ = 456.70 [24160-36-9] ≥99% (HPLC)	5mg 36,000
	NS230602	-20 °C	Dehydrotrametenolic acid [3β-Hydroxylanosta-7,9(11),24-trien-21-oic acid] C ₃₀ H ₄₆ O ₃ = 454.68 [29220-16-4] ≥99% (HPLC)	10mg 50,000
	NS230702	-20 °C	3-O-Acetyl-16α-hydroxytrametenolic acid C ₃₂ H ₅₀ O ₅ = 514.74 [168293-13-8] ≥99% (HPLC)	5mg 50,000
	NS230801	-20 °C	3-O-Acetyl-16α-hydroxydehydrotrametenolic acid C ₃₂ H ₄₈ O ₅ = 512.72 [168293-14-9] ≥99% (HPLC)	2mg 36,000
	NS230901	-20 °C	Polyporenic acid C C ₃₁ H ₄₆ O ₄ = 482.69 [465-18-9] ≥99% (HPLC)	2mg 36,000
	NS231002	-20 °C	Poricoic acid A C ₃₁ H ₄₆ O ₅ = 498.69 [137551-38-3] ≥99% (HPLC)	5mg 36,000
	NS231101	-20 °C	Poricoic acid B C ₃₀ H ₄₄ O ₅ = 484.67 [137551-39-4] ≥99% (HPLC)	2mg 36,000
ウーロン茶 Oolong tea	NS240102	-20 °C	Oolonghomobisflavan A C ₄₅ H ₃₅ O ₂₂ = 928.75 [126737-60-8] ≥98% (HPLC)	5mg 36,000
	NS240202	-20 °C	Oolonghomobisflavan B C ₄₅ H ₃₅ O ₂₂ = 928.75 [176107-91-8] ((126715-88-6)) ≥98% (HPLC)	5mg 36,000
	NS240301	-20 °C	Oolonghomobisflavan C C ₄₅ H ₃₅ O ₂₂ = 928.75 [126716-06-1] ≥98% (HPLC)	2mg 36,000
	NS241101 <i>new</i>	-20 °C	Chafuroside B C ₂₁ H ₁₈ O ₉ = 414.36 [866737-00-0] ≥98% (HPLC)	1mg 40,000
バナバ	NS250102	2-10 °C	Corosolic acid (Colosolic acid) C ₃₀ H ₄₈ O ₄ = 472.70 [4547-24-4] ≥99% (HPLC)	10mg 40,000
	NS250201	2-10 °C	Maslinic acid C ₃₀ H ₄₈ O ₄ = 472.70 [4373-41-5] ≥99% (HPLC)	2mg 20,000
エビ・カニ	NS260102	-20 °C	N-Acetyl-D-glucosamine C ₈ H ₁₅ NO ₆ = 221.21 [7512-17-6] ((72-87-7, 10036-64-3)) ≥α,β Total 99.5% (HPLC), α ≧95%	10mg 30,000
	NS260206	2-10 °C	N-Acetyl-D-glucosamine C ₈ H ₁₅ NO ₆ = 221.21 [7512-17-6] ((72-87-7, 10036-64-3)) ≥α,β Total 99% (HPLC), α ≧95%	100g 12,000
ニンニク、 タマネギ	NS270102	-20 °C	Methiin C ₄ H ₉ NO ₃ S = 151.19 [6853-87-8] ≥99% (HPLC)	1mg 36,000
	NS270202	-20 °C	Propiin C ₆ H ₁₃ NO ₃ S = 179.24 [17795-24-3] ≥99% (HPLC)	1mg 36,000
	NS270302	-20 °C	Isoalliin C ₆ H ₁₁ NO ₃ S = 177.22 [434322-95-9] ≥99% (HPLC)	1mg 36,000
	NS270402	-20 °C	Cycloalliin hydrochloride monohydrate C ₆ H ₁₁ NO ₃ S · HCl · H ₂ O = 231.70 [91673-63-1, Component: 15042-85-0] ≥99% (HPLC)	5mg 36,000
オウゴン 黄連 Coptis japonica	NS280102	2-10 °C	Berberine Chloride C ₂₀ H ₁₈ NO ₄ · Cl = 371.81 [633-65-8, 141433-60-5] ≥99% (HPLC)	20mg 12,000
	NS280202	2-10 °C	Berberrubine Chloride C ₁₉ H ₁₆ NO ₄ · Cl = 357.79 [15401-69-1] ≥99% (HPLC)	10mg 24,000
	NS280302	2-10 °C	Coptisine Chloride C ₁₉ H ₁₄ NO ₄ · Cl = 355.77 [6020-18-4] ≥99% (HPLC)	10mg 30,000
	NS280402	2-10 °C	Epiberberine Chloride C ₂₀ H ₁₈ NO ₄ · Cl = 371.81 [889665-86-5] ≥99% (HPLC)	10mg 36,000
	NS280502	2-10 °C	Palmatine Chloride C ₂₁ H ₂₂ NO ₄ · Cl = 387.86 [10605-02-4] ≥99% (HPLC)	10mg 20,000
	NS280602	2-10 °C	Magnoflorine iodide C ₂₀ H ₂₄ NO ₄ · I = 469.31 [4277-43-4, Component: 2141-09-5] ≥99% (HPLC)	5mg 36,000
	NS281102	2-10 °C	dl-Tetrahydroberberine (dl-Canadine) C ₂₀ H ₂₁ NO ₄ = 339.39 [522-97-4] ≥99% (HPLC)	5mg 40,000
	NS281202	2-10 °C	dl-Tetrahydroberberrubine C ₁₉ H ₁₉ NO ₄ = 325.36 [17388-17-9] ≥99% (HPLC)	5mg 40,000
	NS281302	2-10 °C	dl-Tetrahydrocoptisine (dl-Stylopine) C ₁₉ H ₁₇ NO ₄ = 323.34 [4312-32-7] ≥99% (HPLC)	5mg 40,000
	NS281402	2-10 °C	dl-Tetrahydroepiberberine (dl-Sinactine) C ₂₀ H ₂₁ NO ₄ = 339.39 [38853-67-7] ≥99% (HPLC)	5mg 40,000
	NS281502	2-10 °C	dl-Tetrahydropalmatine C ₂₁ H ₂₅ NO ₄ = 355.43 [2934-97-6] ≥99% (HPLC)	5mg 40,000

Source et. al.	Code No.	Storage Conditions	Product Name・Purity・Molecular Formula= Molecular Weight・〔 CAS Registry Number 〕	Quantity・Price (JPY)
ウメ Mume	NS290102	-20 °C	Mumefural (natural) $\geq 99\%$ (HPLC) 純度規格変更 $C_{12}H_{12}O_9 = 300.22$ [222973-44-6]	5mg 40,000
	NS290202	-20 °C	Mumefural (synthetic) $\geq 99\%$ (HPLC) 純度規格変更 $C_{12}H_{12}O_9 = 300.22$ [222973-44-6]	5mg 40,000
	NS290302	-20 °C	Mumefural isomer (natural) $\geq 99\%$ (HPLC) 純度規格変更 $C_{12}H_{12}O_9 = 300.22$ [1025890-42-9]	2mg 40,000
	NS290402	-20 °C	Mumefural isomer (synthetic) $\geq 98\%$ (HPLC) $C_{12}H_{12}O_9 = 300.22$ [1025890-42-9]	2mg 40,000
ブドウ、 イタドリ、 パッションフルーツ	NS300103	-20 °C	<i>trans</i> -Resveratrol $\geq 99\%$ (HPLC) $C_{14}H_{12}O_3 = 228.24$ [501-36-0]	100mg 10,000
	NS300104	-20 °C	<i>trans</i> -Resveratrol $\geq 99\%$ (HPLC) $C_{14}H_{12}O_3 = 228.24$ [501-36-0]	1g 50,000
	NS300202	-20 °C	<i>cis</i> -Resveratrol $\geq 95\%$ (HPLC) 純度規格変更 $C_{14}H_{12}O_3 = 228.24$ [61434-67-1]	10mg 30,000
ブドウ	NS300602	-20 °C	<i>trans</i> -Pterostilbene $\geq 99\%$ (HPLC) $C_{16}H_{16}O_3 = 256.30$ [537-42-8]	10mg 20,000
	NS300603	-20 °C	<i>trans</i> -Pterostilbene $\geq 99\%$ (HPLC) $C_{16}H_{16}O_3 = 256.30$ [537-42-8]	100mg 140,000
	NS301202	-20 °C	(±)- ϵ -Viniferin (+/- $\approx 1:1$) \geq Total 99% (HPLC) $C_{28}H_{22}O_6 = 454.47$ [253435-07-3]	10mg 20,000
	NS301301	-20 °C	(+)- ϵ -Viniferin $\geq 99\%$ ee (HPLC) $C_{28}H_{22}O_6 = 454.47$ [129170-22-5]	2mg 36,000
	NS301401	-20 °C	(-)- ϵ -Viniferin $\geq 99\%$ ee (HPLC) $C_{28}H_{22}O_6 = 454.47$ [62218-08-0]	2mg 36,000
	NS301501	-20 °C	ω -Viniferin (+/- $\approx 1:3$) \geq Total 98% (HPLC) $C_{28}H_{22}O_6 = 454.47$ [1307823-52-4, 174023-03-1]	2mg 36,000
	NS301602	-20 °C	Ampelopsin A $\geq 99\%$ (HPLC) $C_{28}H_{22}O_7 = 470.47$ [130608-11-6]	5mg 36,000
	NS303101	-20 °C	Miyabenol C $\geq 99\%$ (HPLC) $C_{42}H_{32}O_9 = 680.70$ [109605-83-6]	2mg 50,000
	NS305101	-20 °C	Vitisin A $\geq 99\%$ (HPLC) $C_{56}H_{42}O_{12} = 906.93$ [142449-89-6]	2mg 36,000
	NS305201	-20 °C	Vitisin B $\geq 99\%$ (HPLC) $C_{56}H_{42}O_{12} = 906.93$ [142449-90-9]	2mg 36,000
	NS305301	-20 °C	Hopeaphenol $\geq 99\%$ (HPLC) $C_{56}H_{42}O_{12} = 906.93$ [388582-37-4]	2mg 36,000
	NS305401	-20 °C	Isohopeaphenol $\geq 99\%$ (HPLC) $C_{56}H_{42}O_{12} = 906.93$ [197446-77-8]	2mg 36,000
イタドリ	NS300402	-20 °C	<i>trans</i> -Polydatin (<i>trans</i> -Piceid) $\geq 99\%$ (HPLC) $C_{20}H_{22}O_8 = 390.38$ [27208-80-6]	10mg 20,000
	NS300502	-20 °C	<i>cis</i> -Polydatin (<i>cis</i> -Piceid) $\geq 98\%$ (HPLC) 純度規格変更 $C_{20}H_{22}O_8 = 390.38$ [148766-36-3]	10mg 30,000
パッション フルーツ	NS300302	-20 °C	<i>trans</i> -Piceatannol $\geq 99\%$ (HPLC) $C_{14}H_{12}O_4 = 244.25$ [10083-24-6]	10mg 16,000
	NS300303	-20 °C	<i>trans</i> -Piceatannol $\geq 99\%$ (HPLC) $C_{14}H_{12}O_4 = 244.25$ [10083-24-6]	100mg 128,000
	NS300902	-20 °C	<i>cis</i> -Piceatannol $\geq 95\%$ (HPLC) 純度規格変更 $C_{14}H_{12}O_4 = 244.25$ [106325-86-4]	10mg 36,000
	NS300702	-20 °C	<i>trans</i> -Scirpusin B $\geq 98\%$ (HPLC) $C_{28}H_{22}O_8 = 486.47$ [483363-92-4]	10mg 36,000
	NS301001	-20 °C	<i>cis</i> -Scirpusin B $\geq 98\%$ (HPLC) $C_{28}H_{22}O_8 = 486.47$ [288846-83-3]	2mg 36,000
NS300801	-20 °C	<i>trans</i> -Scirpusin A $\geq 99\%$ (HPLC) $C_{28}H_{22}O_7 = 470.47$ [483363-91-3]	2mg 36,000	
クワ	NS301102	-20 °C	<i>trans</i> -Oxyresveratrol $\geq 99\%$ (HPLC) $C_{14}H_{12}O_4 = 244.24$ [29700-22-9]	10mg 10,000
米糠フェルラ酸	NS310103	2-10 °C	<i>trans</i> -Ferulic acid $\geq 99\%$ (HPLC) $C_{10}H_{10}O_4 = 194.18$ [537-98-4]	100mg 10,000
	NS310104	2-10 °C	<i>trans</i> -Ferulic acid $\geq 99\%$ (HPLC) $C_{10}H_{10}O_4 = 194.18$ [537-98-4]	1g 20,000
キョウカク 羌活 Notopterygium incisum	NS320102	2-10 °C	Nodakenin $\geq 99\%$ (HPLC) $C_{20}H_{24}O_9 = 408.40$ [495-31-8]	5mg 36,000
	NS320201	2-10 °C	Marmesinin $\geq 99\%$ (HPLC) $C_{20}H_{24}O_9 = 408.40$ [495-30-7]	2mg 36,000
	NS320302	2-10 °C	Notopterol $\geq 99\%$ (HPLC) $C_{21}H_{22}O_5 = 354.40$ [88206-46-6]	10mg 30,000
	NS320402	2-10 °C	Notoptol $\geq 99\%$ (HPLC) $C_{21}H_{22}O_5 = 354.40$ [88206-49-9]	5mg 36,000
ダイオウ 大黃 Rheum palmatum	NS330102	2-10 °C	Rhein $\geq 99\%$ (HPLC) $C_{15}H_8O_6 = 284.22$ [478-43-3]	20mg 20,000
NS330103	2-10 °C	Rhein $\geq 99\%$ (HPLC) $C_{15}H_8O_6 = 284.22$ [478-43-3]	100mg 80,000	

Source et. al.	Code No.	Storage Conditions	Product Name・Purity・Molecular Formula=Molecular Weight・〔 CAS Registry Number 〕	Quantity・Price (JPY)
カシス、黒大豆他	NS380502	-20 °C	Cyanidin chloride $\geq 99\%$ (HPLC) C ₁₅ H ₁₁ O ₆ ・Cl=322.70 [528-58-5, Component: 13306-05-3]	5mg 10,000
	NS380101 NS380102	-20 °C	Cyanidin 3-glucoside chloride [Kuromanin chloride] $\geq 99\%$ (HPLC) C ₂₁ H ₂₁ O ₁₁ ・Cl=484.84 [7084-24-4, Component: 47705-70-4]	5mg 15,000 10mg 26,000
	NS380202	-20 °C	Cyanidin 3-rutinoside chloride [Keracyanin chloride] $\geq 99\%$ (HPLC) C ₂₇ H ₃₁ O ₁₅ ・Cl=630.98 [18719-76-1, Component: 28338-59-2]	10mg 12,000
	NS380602	-20 °C	Delphinidin chloride $\geq 99\%$ (HPLC) C ₁₅ H ₁₁ O ₇ ・Cl=338.70 [528-53-0, Component: 13270-61-6]	5mg 16,000
	NS380301 NS380302	-20 °C	Delphinidin 3-glucoside chloride [Mirtillin chloride] $\geq 99\%$ (HPLC) C ₂₁ H ₂₁ O ₁₂ ・Cl=500.84 [6906-38-3, Component: 50986-17-9]	5mg 20,000 10mg 36,000
	NS380401 NS380402	-20 °C	Delphinidin 3-rutinoside chloride [Tulipanin chloride] $\geq 99\%$ (HPLC) C ₂₇ H ₃₁ O ₁₆ ・Cl=646.98 [15674-58-5, Component: 58285-26-0]	5mg 24,000 10mg 36,000
イヌリン Inulin	NS390102	2-10 °C	Fructo-oligosaccharide DP5 [GF ₄ , Fructofuranosyl-nystose] Fru β (2-1)-[Fru β (2-1)] ₃ - α (2-1)Glc $\geq 99\%$ (HPLC) C ₃₀ H ₅₂ O ₂₆ = 828.71 []	5mg 32,000
	NS390201	2-10 °C	Fructo-oligosaccharide DP6 [GF ₅ , 1-F-(1- β -fructofuranosyl)-2-nystose] Fru β (2-1)-[Fru β (2-1)] ₄ - α (2-1)Glc $\geq 99\%$ (HPLC) C ₃₆ H ₆₂ O ₃₁ = 990.85 []	1mg 32,000
	NS390301	2-10 °C	Fructo-oligosaccharide DP7 [GF ₆ , 1-F-(1- β -fructofuranosyl)-3-nystose] Fru β (2-1)-[Fru β (2-1)] ₅ - α (2-1)Glc $\geq 99\%$ (HPLC) C ₄₂ H ₇₂ O ₃₆ = 1152.99 []	1mg 32,000
	NS390401	2-10 °C	Fructo-oligosaccharide DP8 [GF ₇] Fru β (2-1)-[Fru β (2-1)] ₆ - α (2-1)Glc $\geq 99\%$ (HPLC) C ₄₈ H ₈₂ O ₄₁ = 1315.13 []	1mg 32,000
	NS390501	2-10 °C	Fructo-oligosaccharide DP9 [GF ₈] Fru β (2-1)-[Fru β (2-1)] ₇ - α (2-1)Glc $\geq 99\%$ (HPLC) C ₅₄ H ₉₂ O ₄₆ = 1477.27 []	1mg 32,000
	NS390601	2-10 °C	Fructo-oligosaccharide DP10 [GF ₉] Fru β (2-1)-[Fru β (2-1)] ₈ - α (2-1)Glc $\geq 99\%$ (HPLC) C ₆₀ H ₁₀₂ O ₅₁ = 1639.41 []	1mg 32,000
	NS390701	2-10 °C	Fructo-oligosaccharide DP11 [GF ₁₀] Fru β (2-1)-[Fru β (2-1)] ₉ - α (2-1)Glc $\geq 99\%$ (HPLC) C ₆₆ H ₁₁₂ O ₅₆ = 1801.55 []	1mg 32,000
	NS390801	2-10 °C	Fructo-oligosaccharide DP12 [GF ₁₁] Fru β (2-1)-[Fru β (2-1)] ₁₀ - α (2-1)Glc $\geq 99\%$ (HPLC) C ₇₂ H ₁₂₂ O ₆₁ = 1963.69 []	1mg 32,000
	NS390901	2-10 °C	Fructo-oligosaccharide DP13 [GF ₁₂] Fru β (2-1)-[Fru β (2-1)] ₁₁ - α (2-1)Glc $\geq 99\%$ (HPLC) C ₇₈ H ₁₃₂ O ₆₆ = 2125.83 []	1mg 32,000
	NS391001	2-10 °C	Fructo-oligosaccharide DP14 [GF ₁₃] Fru β (2-1)-[Fru β (2-1)] ₁₂ - α (2-1)Glc $\geq 99\%$ (HPLC) C ₈₄ H ₁₄₂ O ₇₁ = 2287.97 []	1mg 32,000
	NS391101	2-10 °C	Fructo-oligosaccharide DP15 [GF ₁₄] Fru β (2-1)-[Fru β (2-1)] ₁₃ - α (2-1)Glc $\geq 99\%$ (HPLC) C ₉₀ H ₁₅₂ O ₇₆ = 2450.11 []	1mg 32,000
	NS391201	2-10 °C	Fructo-oligosaccharide DP16 [GF ₁₅] Fru β (2-1)-[Fru β (2-1)] ₁₄ - α (2-1)Glc $\geq 99\%$ (HPLC) C ₉₆ H ₁₆₂ O ₈₁ = 2612.25 []	1mg 32,000
	NS391301	2-10 °C	Fructo-oligosaccharide DP17 [GF ₁₆] Fru β (2-1)-[Fru β (2-1)] ₁₅ - α (2-1)Glc $\geq 99\%$ (HPLC) C ₁₀₂ H ₁₇₂ O ₈₆ = 2774.39 []	1mg 32,000
	NS391401	2-10 °C	Fructo-oligosaccharide DP18 [GF ₁₇] Fru β (2-1)-[Fru β (2-1)] ₁₆ - α (2-1)Glc $\geq 95\%$ (HPLC) C ₁₀₈ H ₁₈₂ O ₉₁ = 2936.53 []	0.2mg 32,000
	NS391501	2-10 °C	Fructo-oligosaccharide DP19 [GF ₁₈] Fru β (2-1)-[Fru β (2-1)] ₁₇ - α (2-1)Glc $\geq 95\%$ (HPLC) C ₁₁₄ H ₁₉₂ O ₉₆ = 3098.67 []	0.2mg 32,000
	NS391601	2-10 °C	Fructo-oligosaccharide DP20 [GF ₁₉] Fru β (2-1)-[Fru β (2-1)] ₁₈ - α (2-1)Glc $\geq 95\%$ (HPLC) C ₁₂₀ H ₂₀₂ O ₁₀₁ = 3260.81 []	0.2mg 32,000
	NS395101	2-10 °C	Fructo-oligosaccharides kit (DP3~DP11, GF ₂ ~GF ₁₀)	1mg 1set 200,000
	NS395201	2-10 °C	Fructo-oligosaccharides kit (DP12~DP20, GF ₁₁ ~GF ₁₉)	DP12~DP17, 1mg DP18~DP20, 0.2mg 1set 200,000

Source et. al.	Code No.	Storage Conditions	Product Name · Molecular Formula = Molecular Weight · [CAS Registry Number]	Purity	Quantity · Price (JPY)
希少糖 Rare sugar (Keto-hexoses)	NS400103	2-10 °C	D-Psicose C ₆ H ₁₂ O ₆ =180.16 [551-68-8]	≥99.5% (HPLC)	100mg 12,000
	NS400204	2-10 °C		≥99% (HPLC)	1g 32,000
	NS400303	2-10 °C	L-Psicose C ₆ H ₁₂ O ₆ =180.16 [16354-64-6]	≥99.5% (HPLC)	100mg 36,000
	NS400403	2-10 °C		≥99% (HPLC)	100mg 12,000
	NS400503	2-10 °C	L-Fructose C ₆ H ₁₂ O ₆ =180.16 [7776-48-9]	—	—
	NS400604	2-10 °C		≥98% (HPLC)	1g 42,000
	NS400703	2-10 °C	D-Sorbose C ₆ H ₁₂ O ₆ =180.16 [3615-56-3]	≥99.5% (HPLC)	100mg 12,000
	NS400804	2-10 °C		≥99% (HPLC)	1g 15,000
	NS400903	2-10 °C	L-Sorbose C ₆ H ₁₂ O ₆ =180.16 [87-79-6]	≥99.5% (HPLC)	100mg 12,000
	NS401005	2-10 °C		≥99% (HPLC)	25g 10,000
	NS401103	2-10 °C	D-Tagatose C ₆ H ₁₂ O ₆ =180.16 [87-81-0]	≥99.5% (HPLC)	100mg 12,000
	NS401204	2-10 °C		≥98% (HPLC)	1g 10,000
	NS401303	2-10 °C	L-Tagatose C ₆ H ₁₂ O ₆ =180.16 [17598-82-2]	≥99.5% (HPLC)	100mg 60,000
	NS401403	2-10 °C		≥98% (HPLC)	100mg 20,000
希少糖 Rare sugar (Aldo-hexoses)	NS401503	2-10 °C	D-Allose C ₆ H ₁₂ O ₆ =180.16 [2595-97-3]	≥99.5% (HPLC)	100mg 12,000
	NS401604	2-10 °C		≥99% (HPLC)	1g 15,000
	NS401703	2-10 °C	D-Altrose C ₆ H ₁₂ O ₆ =180.16 [1990-29-0]	≥99.5% (HPLC)	100mg 15,000
	NS401804	2-10 °C		≥99% (HPLC)	1g 32,000
希少糖 Rare sugar (Sugar alcohols)	NS401903	2-10 °C	Allitol C ₆ H ₁₄ O ₆ =182.17 [488-44-8]	≥99.5% (HPLC)	100mg 12,000
	NS402004	2-10 °C		≥99% (HPLC)	1g 30,000
	NS402103	2-10 °C	D-Iditol C ₆ H ₁₄ O ₆ =182.17 [25878-23-3]	≥99.5% (HPLC)	100mg 20,000
	NS402204	2-10 °C		≥99% (HPLC)	1g 38,000
	NS402303	2-10 °C	L-Iditol C ₆ H ₁₄ O ₆ =182.17 [488-45-9]	≥99.5% (HPLC)	100mg 25,000
	NS402404	2-10 °C		≥99% (HPLC)	1g 45,000
	NS402503	2-10 °C	D-Talitol C ₆ H ₁₄ O ₆ =182.17 [643-03-8]	≥99.5% (HPLC)	100mg 30,000
	NS402604	2-10 °C		≥99% (HPLC)	0.5g 30,000
	NS402703	2-10 °C	L-Talitol C ₆ H ₁₄ O ₆ =182.17 [60660-58-4]	≥99.5% (HPLC)	100mg 75,000
	NS402803	2-10 °C		≥99% (HPLC)	200mg 36,000
	NS402903	2-10 °C	L-Sorbitol C ₆ H ₁₄ O ₆ =182.17 [6706-59-8]	≥99.5% (HPLC)	100mg 50,000
	NS403003	2-10 °C		≥99% (HPLC)	100mg 25,000
	NS403103	2-10 °C	L-Mannitol C ₆ H ₁₄ O ₆ =182.17 [643-01-6]	≥99.5% (HPLC)	100mg 12,000
	NS403204	2-10 °C		≥99% (HPLC)	1g 20,000

Source et. al.	Code No.	Storage Conditions	Product Name · Purity · Molecular Formula = Molecular Weight · [CAS Registry Number]	Quantity · Price (JPY)
高純度飽和脂肪酸 Saturated fatty acids	NS410105	2-10 °C	Decanoic acid [Capric acid] $\geq 99.5\%$ (HPLC) $C_{10}H_{20}O_2=172.26$ [334-48-5]	10g 12,000
	NS410205	2-10 °C	Dodecanoic acid [Lauric acid] $\geq 99.5\%$ (HPLC) $C_{12}H_{24}O_2=200.32$ [143-07-7]	10g 12,000
	NS411105	2-10 °C	Tridecanoic acid $\geq 99.5\%$ (HPLC) $C_{13}H_{26}O_2=214.35$ [638-53-9]	10g 12,000
	NS410305	2-10 °C	Myristic acid $\geq 99.5\%$ (HPLC) $C_{14}H_{28}O_2=228.37$ [544-63-8]	10g 12,000
	NS411205	2-10 °C	Pentadecanoic acid $\geq 99.5\%$ (HPLC) $C_{15}H_{30}O_2=242.40$ [1002-84-2]	10g 12,000
	NS410405	2-10 °C	Palmitic acid $\geq 99.5\%$ (HPLC) $C_{16}H_{32}O_2=256.42$ [57-10-3]	10g 12,000
	NS411305	2-10 °C	Heptadecanoic acid $\geq 99.5\%$ (HPLC) $C_{17}H_{34}O_2=270.46$ [506-12-7]	5g 12,000
	NS410505	2-10 °C	Stearic acid $\geq 99.5\%$ (HPLC) $C_{18}H_{36}O_2=284.48$ [57-11-4]	10g 12,000
	NS411404	2-10 °C	Nonadecanoic acid $\geq 99.5\%$ (HPLC) $C_{19}H_{38}O_2=298.51$ [646-30-0]	2g 12,000
	NS410604	2-10 °C	Arachidic acid $\geq 99.5\%$ (HPLC) $C_{20}H_{40}O_2=312.53$ [506-30-9]	1g 12,000
	NS411503	2-10 °C	Heneicosanoic acid $\geq 99.5\%$ (HPLC) $C_{21}H_{42}O_2=326.57$ [2363-71-5]	200mg 12,000
	NS410704	2-10 °C	Behenic acid $\geq 99.5\%$ (HPLC) $C_{22}H_{44}O_2=340.58$ [112-85-6]	1g 12,000
	NS411603	2-10 °C	Tricosanoic acid $\geq 99.5\%$ (HPLC) $C_{23}H_{46}O_2=354.62$ [2433-96-7]	100mg 12,000
	NS410804	2-10 °C	Lignoceric acid $\geq 99.5\%$ (HPLC) $C_{24}H_{48}O_2=368.63$ [557-59-5]	1g 15,000
	NS411703	2-10 °C	Pentacosanoic acid $\geq 99.5\%$ (HPLC) $C_{25}H_{50}O_2=382.67$ [506-38-7]	100mg 20,000
	NS410904	2-10 °C	Hexacosanoic acid $\geq 99.5\%$ (HPLC) $C_{26}H_{52}O_2=396.69$ [506-46-7]	500mg 20,000
	NS411803	2-10 °C	Heptacosanoic acid $\geq 99.5\%$ (TLC) $C_{27}H_{54}O_2=410.73$ [7138-40-1]	100mg 20,000
	NS411003	2-10 °C	Octacosanoic acid $\geq 99.5\%$ (HPLC) $C_{28}H_{56}O_2=424.75$ [506-48-9]	100mg 36,000
	NS411903	2-10 °C	Nonacosanoic acid $\geq 99.5\%$ (TLC) $C_{29}H_{58}O_2=438.78$ [4250-38-8]	100mg 36,000
	NS412003	2-10 °C	Melissic acid $\geq 99.5\%$ (TLC) $C_{30}H_{60}O_2=452.81$ [506-50-3]	100mg 36,000
高純度飽和脂肪酸 銀塩	NS470102	冷暗所	Silver behenate $\geq 99.5\%$ $C_{22}H_{43}AgO_2=447.44$ [2489-05-6]	5mg 36,000
ブロッコリー Broccoli	NS420102	-20 °C	Glucoraphanin $\geq 98\%$ (HPLC)	10mg 28,000
	NS420103	-20 °C	$C_{12}H_{23}NO_{10}S_3 = 437.51$ [21414-41-5]	100mg 224,000
	NS420201	-20 °C	Glucobrerin $\geq 98\%$ (HPLC) $C_{11}H_{21}NO_{10}S_3 = 423.48$ [554-88-1]	2mg 28,000
コーヒー Coffee	NS430102	-20 °C	3-O-Caffeoylquinic acid [Neochlorogenic acid] $\geq 98\%$ (HPLC) $C_{16}H_{18}O_9 = 354.31$ [342811-68-1, 906-33-2] *注	5mg 30,000
	NS430202	-20 °C	4-O-Caffeoylquinic acid $\geq 98\%$ (HPLC) $C_{16}H_{18}O_9 = 354.31$ [905-99-7, 87099-73-8]	5mg 30,000
	NS430302	-20 °C	Chlorogenic acid [5-O-Caffeoylquinic acid] $\geq 99\%$ (HPLC) $C_{16}H_{18}O_9 = 354.31$ [327-97-9, 202650-88-2] *注	10mg 24,000
	NS430401	-20 °C	3-O-Feruloylquinic acid $\geq 98\%$ (HPLC) $C_{17}H_{20}O_9 = 368.34$ [40242-06-6, 87099-72-7] *注	1mg 45,000
	NS430501	-20 °C	4-O-Feruloylquinic acid $\geq 98\%$ (HPLC) $C_{17}H_{20}O_9 = 368.34$ [2613-86-7, 96646-16-1]	1mg 45,000
	NS430602	-20 °C	5-O-Feruloylquinic acid $\geq 98\%$ (HPLC) $C_{17}H_{20}O_9 = 368.34$ [53905-80-9, 62929-69-5] *注	5mg 30,000
防己	NS450102	2-10 °C	Tetrandrine $\geq 99.5\%$ (HPLC)	10mg 12,000
	NS450103	2-10 °C	$C_{28}H_{42}N_2O_6 = 622.75$ [518-34-3]	100mg 36,000
	NS450202	2-10 °C	Fangchinoline $\geq 99\%$ (HPLC) $C_{37}H_{40}N_2O_6 = 608.72$ [436-77-1]	5mg 30,000

*注：化合物名はM. Murata, H.Okada, and S. Homma, *Biosci. Biotech. Biochem.*, **59** (10), 1887-1890 (1995). による。

Source et. al.	Code No.	Storage Conditions	Product Name · Purity · Molecular Formula = Molecular Weight · [CAS Registry Number]	Quantity · Price (JPY)
グルコシルセラミド等 Glucosylceramides etc.	NS170102 NS170103	2-10 °C	Glucosylceramide, from Rice (Glucosylceramide mix.) (米) [85305-87-9] $\geq 99\%$ (TLC)	10mg 16,000 100mg 110,000
	NS170202 NS170203	2-10 °C	Glucosylceramide, from Maize (Glucosylceramide mix.) (トウモロコシ) [85305-87-9] $\geq 99\%$ (TLC)	10mg 15,000 100mg 110,000
	NS170302 NS170303	2-10 °C	Glucosylceramide, from Konjac (Glucosylceramide mix.) (蒟蒻) [85305-87-9] $\geq 99\%$ (TLC)	10mg 20,000 100mg 150,000
	NS170402 NS170403	2-10 °C	Glucosylceramide, from Hen of the Woods (Maitake) (Glucosylceramide mix.) (舞茸) [85305-87-9] $\geq 99\%$ (TLC)	10mg 20,000 100mg 150,000
	NS170502 NS170503	2-10 °C	Glucosylceramide, from Tamogitake (Glucosylceramide mix.) (たもぎ茸) [85305-87-9] $\geq 99\%$ (TLC)	10mg 20,000 100mg 150,000
	NS170602 NS170603	2-10 °C	Glucosylceramide, from Soybean (Glucosylceramide mix.) (大豆) [497155-61-0] $\geq 99\%$ (TLC)	10mg 16,000 100mg 110,000
	NS170702 NS170703	2-10 °C	Glucosylceramide, from Wheat (Glucosylceramide mix.) (小麦) [85305-87-9] $\geq 99\%$ (TLC)	10mg 20,000 100mg 150,000
	NS170802 NS170803	2-10 °C	Glucosylceramide, from Sugar beet (Glucosylceramide mix.) (ビート) [85305-87-9] (学名: Beta vulgaris L. subsp. vulgaris var. altissima) $\geq 99\%$ (TLC)	10mg 20,000 100mg 150,000
	NS170902 NS170903	2-10 °C	Ceramide, from <i>Acetobacter malorum</i> (Ceramide mix.) (酢酸菌) [100403-19-8] $\geq 99\%$ (TLC)	10mg 20,000 100mg 150,000
	NS220302 NS220303	2-10 °C	Glucosylceramide, from Milk (Glucosylceramide mix.) (ミルク) [85305-87-9] $\geq 99\%$ (TLC)	10mg 20,000 100mg 150,000
	NS220402 NS220403	2-10 °C	Lactosylceramide, from Milk (Lactosylceramide mix.) (ミルク) [4682-48-8] $\geq 99\%$ (TLC)	10mg 20,000 100mg 150,000
	NS171002 NS171003	2-10 °C	Glucosylceramide, from Peach (Glucosylceramide mix.) (桃) [85305-87-9] $\geq 99\%$ (TLC)	10mg 20,000 100mg 150,000
	NS171102	2-10 °C	Ceramide, from Citrus unshiu (Ceramide mix.) (みかん) [100403-19-8] $\geq 99\%$ (TLC)	5mg 40,000
	NS171202	2-10 °C	Glucosylceramide, from Citrus unshiu (Glucosylceramide mix.) (みかん) [85305-87-9] $\geq 99\%$ (TLC)	5mg 60,000
	NS171302 <i>new</i>	2-10 °C	Glucosylceramide, from <i>Aspergillus oryzae</i> (Glucosylceramide mix.) (麹菌) [] $\geq 99\%$ (TLC)	5mg 30,000
	NS171402 <i>new</i>	2-10 °C	Glucosylceramide, from Red beet (Glucosylceramide mix.) (ビーツ) [85305-87-9] (学名: Beta vulgaris L. subsp. vulgaris var. vulgaris) $\geq 99\%$ (TLC)	5mg 45,000
グルコシルセラミド構成成分 ・米 Glucosylceramide, from Rice	NS340101	2-10 °C	d18:2 ^{4E,8Z} -C20h:0 Glucosylceramide, from Rice C ₄₄ H ₈₃ NO ₉ = 770.13 [] $\geq 99\%$ (HPLC)	2mg 40,000
	NS340201	2-10 °C	d18:2 ^{4E,8Z} -C20h:0 Glucosylceramide, from Rice C ₄₄ H ₈₃ NO ₉ = 770.13 [] $\geq 99\%$ (HPLC)	1mg 40,000
	NS340301	2-10 °C	t18:1 ^{8Z} -C22h:0 Glucosylceramide, from Rice C ₄₆ H ₈₉ NO ₁₀ = 816.20 [] $\geq 99\%$ (HPLC)	1mg 40,000
	NS340401	2-10 °C	d18:2 ^{4E,8Z} -C22h:0 Glucosylceramide, from Rice C ₄₆ H ₈₇ NO ₉ = 798.18 [] $\geq 99\%$ (HPLC)	1mg 60,000
	NS340501	2-10 °C	t18:1 ^{8Z} -C24h:0 Glucosylceramide, from Rice C ₄₈ H ₉₃ NO ₁₀ = 844.25 [] $\geq 99\%$ (HPLC)	1mg 40,000
	NS340601	2-10 °C	d18:2 ^{4E,8Z} -C24h:0 Glucosylceramide, from Rice C ₄₈ H ₉₁ NO ₉ = 826.24 [] $\geq 99\%$ (HPLC)	1mg 60,000
グルコシルセラミド構成成分 ・蒟蒻 Glucosylceramide, from Konjac	NS350101	2-10 °C	d18:2 ^{4E,8Z} -C16h:0 Glucosylceramide, from Konjac C ₄₀ H ₇₅ NO ₉ = 714.02 [] $\geq 99\%$ (HPLC)	2mg 40,000
	NS350201	2-10 °C	d18:2 ^{4E,8Z} -C16h:0 Glucosylceramide, from Konjac C ₄₀ H ₇₅ NO ₉ = 714.02 [] $\geq 99\%$ (HPLC)	1mg 40,000
	NS350301	2-10 °C	d18:2 ^{4E,8Z} -C18h:0 Glucosylceramide, from Konjac C ₄₂ H ₇₉ NO ₉ = 742.08 [] $\geq 99\%$ (HPLC)	2mg 40,000
	NS350401	2-10 °C	d18:2 ^{4E,8Z} -C18h:0 Glucosylceramide, from Konjac C ₄₂ H ₇₉ NO ₉ = 742.08 [] $\geq 99\%$ (HPLC)	1mg 40,000
	NS350501	2-10 °C	d18:2 ^{4E,8Z} -C20h:0 Glucosylceramide, from Konjac C ₄₄ H ₈₃ NO ₉ = 770.13 [] $\geq 99\%$ (HPLC)	1mg 40,000
	NS350601	2-10 °C	d18:2 ^{4E,8Z} -C20h:0 Glucosylceramide, from Konjac C ₄₄ H ₈₃ NO ₉ = 770.13 [] $\geq 99\%$ (HPLC)	1mg 60,000
	NS350701	2-10 °C	t18:1 ^{8Z} -C22h:0 Glucosylceramide, from Konjac C ₄₆ H ₈₉ NO ₁₀ = 816.20 [] $\geq 99\%$ (HPLC)	2mg 40,000
	NS350801	2-10 °C	t18:1 ^{8Z} -C23h:0 Glucosylceramide, from Konjac C ₄₇ H ₉₁ NO ₁₀ = 830.23 [] $\geq 99\%$ (HPLC)	0.1mg 40,000
	NS350901	2-10 °C	t18:1 ^{8Z} -C24h:0 Glucosylceramide, from Konjac C ₄₈ H ₉₃ NO ₁₀ = 844.25 [] $\geq 99\%$ (HPLC)	1mg 40,000

Source et. al.	Code No.	Storage Conditions	Product Name · Purity · Molecular Formula = Molecular Weight · [CAS Registry Number]	Quantity · Price (JPY)			
グルコシルセラミド構成成分・舞茸 Glucosylceramide, from Hen of the Woods (Maitake)	NS360101	2-10 °C	9-Me d18:2 ^{4E,8E} -C14h:0 Glucosylceramide, from Hen of the Woods (Maitake) C ₃₉ H ₇₃ NO ₉ = 700.00 [] ≥99% (HPLC)	要問合せ			
	NS360201	2-10 °C	9-Me d18:2 ^{4E,8E} -C15h:0 Glucosylceramide, from Hen of the Woods (Maitake) C ₄₀ H ₇₅ NO ₉ = 714.02 [] ≥99% (HPLC)	1mg 40,000			
	NS360302	2-10 °C	9-Me d18:2 ^{4E,8E} -C16h:0 Glucosylceramide, from Hen of the Woods (Maitake) C ₄₁ H ₇₇ NO ₉ = 728.05 [] ≥99% (HPLC)	5mg 40,000			
	NS360401	2-10 °C	9-Me d18:2 ^{4E,8E} -C17h:0 Glucosylceramide, from Hen of the Woods (Maitake) C ₄₂ H ₇₉ NO ₉ = 742.08 [] ≥99% (HPLC)	1mg 40,000			
	NS360501	2-10 °C	9-Me d18:2 ^{4E,8E} -C18h:0 Glucosylceramide, from Hen of the Woods (Maitake) C ₄₃ H ₈₁ NO ₉ = 756.10 [] ≥99% (HPLC)	1mg 40,000			
グルコシルセラミド構成成分・たもぎ茸 Glucosylceramide, from Tamogitake	NS370101	2-10 °C	9-Me d18:2 ^{4E,8E} -C15h:0 Glucosylceramide, from Tamogitake ≥99% (HPLC) C ₄₀ H ₇₅ NO ₉ = 714.02 []	要問合せ			
	NS370202	2-10 °C	9-Me d18:2 ^{4E,8E} -C16h:0 Glucosylceramide, from Tamogitake ≥99% (HPLC) C ₄₁ H ₇₇ NO ₉ = 728.05 []	5mg 40,000			
	NS370301	2-10 °C	9-Me d18:2 ^{4E,8E} -C17h:0 Glucosylceramide, from Tamogitake ≥99% (HPLC) C ₄₂ H ₇₉ NO ₉ = 742.08 []	要問合せ			
	NS370401	2-10 °C	9-Me d18:2 ^{4E,8E} -C18h:0 Glucosylceramide, from Tamogitake ≥99% (HPLC) C ₄₃ H ₈₁ NO ₉ = 756.10 []	1mg 40,000			
リン脂質・ミルク Phospholipids, from Milk	NS220102	-20°C	Sphingomyelin, from Milk (Sphingomyelin mix.) [475662-40-9]	≥99%(TLC)	20mg 10,000		
	NS220103			100mg 20,000			
	NS220204	-20°C		≥98%(TLC)	1g 24,000		
	NS220702	-20°C	Phosphatidylcholine, from Milk (Phosphatidylcholine mix.) [8002-43-5]	≥98%(TLC)	10mg 20,000		
	NS220802	-20°C	Phosphatidylethanolamine, from Milk (Phosphatidylethanolamine mix.)	≥98%(TLC)	10mg 24,000		
	NS220902	-20°C	Phosphatidylinositol, from Milk (Phosphatidylinositol mix.)	≥98%(TLC)	10mg 36,000		
	NS221002	-20°C	Phosphatidylserine, from Milk (Phosphatidylserine mix.) [1446756-47-3]	≥98%(TLC)	10mg 36,000		
NS221102	-20°C	Lactosyl phosphatidylethanolamine, from Milk (Lactosyl phosphatidylethanolamine mix.)	≥98%(TLC)	5mg 48,000			
リン脂質・卵黄 Phospholipids, from Egg yolk	NS210102	-20°C	Sphingomyelin, from Egg yolk (Sphingomyelin mix.) [383907-87-7]	≥99%(TLC)	20mg 10,000		
	NS210103			100mg 20,000			
	NS210204	-20°C		≥98%(TLC)	終売いたしました		
ガングリオシド・ミルク Gangliosides, from Milk	NS220502	-20°C	Ganglioside GM3, from Milk [54827-14-4]	≥99%(TLC)	5mg 36,000		
	NS220602	-20°C	Ganglioside GD3, ammonium salt, from Milk	≥99%(TLC)	5mg 36,000		
ガングリオシド・合成 Gangliosides (synthetic) etc.	GM200101	-20°C	Ganglioside GM2 (18, 18) C ₆₇ H ₁₂₁ N ₃ O ₂₅ = 1384.68 [127663-77-8]	≥99%(TLC)	0.2mg 50,000		
	GM300101	-20°C	Ganglioside GM3 (18, 2) C ₄₃ H ₇₃ N ₂ O ₂₁ = 957.06 [111545-33-6]	≥99%(TLC)	0.2mg 30,000		
	GM300201	-20°C	Ganglioside GM3 (18, 8) C ₄₉ H ₈₃ N ₂ O ₂₁ = 1041.22 []	≥99%(TLC)	0.2mg 30,000		
	GM300301	-20°C	Ganglioside GM3 (18, 12) C ₅₃ H ₉₃ N ₂ O ₂₁ = 1097.33 [1448012-42-7]	≥99%(TLC)	0.2mg 30,000		
	GM300401	-20°C	Ganglioside GM3 (18, 16) C ₅₇ H ₁₀₃ N ₂ O ₂₁ = 1153.44 [138749-26-5]	≥99%(TLC)	0.2mg 30,000		
	GM300501	-20°C	Ganglioside GM3 (18, 18) C ₆₀ H ₁₀₈ N ₂ O ₂₁ = 1181.49 [124579-05-1]	≥99%(TLC)	0.2mg 30,000		
	GM300601	-20°C	Ganglioside GM3 (18, 24) C ₆₅ H ₁₂₀ N ₂ O ₂₁ = 1265.65 [98461-64-4]	≥99%(TLC)	0.2mg 30,000		
	GX001010 GX001100 GX001500	-20°C	Azido-erythro-sphingosine C ₁₈ H ₃₅ N ₃ O ₂ = 325.49 [103348-49-8]	≥99%(TLC)	50mg 25,000 100mg 40,000 500mg 160,000		
	GX002010 GX002100			-20°C	(2 <i>S</i> , 3 <i>R</i> , 4 <i>E</i>)-2-Azido-3-benzoyl-erythro-sphingosine C ₂₅ H ₃₉ N ₃ O ₃ = 429.60 [103348-50-1]	≥99%(TLC)	10mg 23,000 100mg 180,000
	GX003010 GX003100			-20°C	(2 <i>S</i> , 3 <i>R</i> , 4 <i>E</i>)-2-Azido-3-(<i>tert</i> -butyldimethylsilyl)-erythro-sphingosine ≥99%(TLC) C ₂₄ H ₄₉ N ₃ O ₂ Si = 439.75 [114299-64-8]		10mg 23,000 100mg 180,000

Source et. al.	Code No.	Storage Conditions	Product Name · Purity · Molecular Formula = Molecular Weight · [CAS Registry Number]	Quantity · Price (JPY)
スフィンゴイド塩基 Sphingoid bases	NS440101 NS440102	-20 °C	<i>trans</i> -4- <i>cis</i> -8-Sphingadienine C ₁₈ H ₃₅ NO ₂ = 297.48 [41679-33-8]	≥98% (HPLC) 1mg 40,000 5mg 120,000
	NS440801 <i>new</i>	-20 °C	<i>trans</i> -4- <i>trans</i> -8-Sphingadienine C ₁₈ H ₃₅ NO ₂ = 297.48 [40878-79-3]	≥98% (HPLC) 1mg 50,000
	NS440201 NS440202	-20 °C	4-Hydroxy- <i>cis</i> -8-sphingenine C ₁₈ H ₃₇ NO ₃ = 315.49 [218786-92-6]	≥98% (HPLC) 1mg 65,000 5mg 200,000
	NS440901 <i>new</i>	-20 °C	9-Methyl- <i>trans</i> -4- <i>trans</i> -8-sphingadienine C ₁₉ H ₃₇ NO ₂ = 311.50 [160226-28-8]	≥98% (HPLC) 1mg 60,000
	NS440302	-20 °C	C ₁₂ -Sphingosine C ₁₂ H ₂₅ NO ₂ = 215.33 [128427-86-1]	≥99% (HPLC) 10mg 36,000
	NS440402	-20 °C	C ₁₄ -Sphingosine C ₁₄ H ₂₉ NO ₂ = 243.39 [24558-60-9]	≥99% (HPLC) 10mg 36,000
	NS440502	-20 °C	C ₁₆ -Sphingosine C ₁₆ H ₃₃ NO ₂ = 271.44 [6982-09-8]	≥99% (HPLC) 10mg 36,000
	NS440602	-20 °C	C ₁₈ -Sphingosine C ₁₈ H ₃₇ NO ₂ = 299.49 [123-78-4]	≥99% (HPLC) 25mg 14,000
	NS440702	-20 °C	C ₂₀ -Sphingosine C ₂₀ H ₄₁ NO ₂ = 327.55 [6918-49-6]	≥99% (HPLC) 10mg 36,000
ラノリン・ステロイド	NS460102 NS460103	-20 °C	Lanosterol C ₃₀ H ₅₀ O = 426.72 [79-63-0]	≥99.5% (HPLC) 10mg 20,000 100mg 120,000
	NS460201 NS460202	-20 °C	24, 25-Dihydrolanosterol C ₃₀ H ₅₂ O = 428.73 [79-62-9]	≥99.5% (HPLC) 5mg 20,000 10mg 36,000
	NS460303 NS460304	-20 °C	Cholesterol C ₂₇ H ₄₆ O = 386.65 [57-88-5]	≥99.5% (HPLC) 100mg 15,000 1g 24,000
	NS460402	-20 °C	Desmosterol [24-Dehydrocholesterol] C ₂₇ H ₄₄ O = 384.64 [313-04-2]	≥99% (HPLC) 10mg 36,000
	NS460502	-20 °C	Lathosterol C ₂₇ H ₄₆ O = 386.65 [80-99-9]	≥99% (HPLC) 10mg 36,000
アンテナ型 ラノリン脂肪酸	NS480101	2-10 °C	18-Methyleicosanoic acid [18-MEA] (natural) C ₂₁ H ₄₂ O ₂ = 326.56 [340257-50-3] ([36332-93-1])	≥99% (HPLC) 1mg 36,000
	NS480201	2-10 °C	20-Methyldocosanoic acid (natural) C ₂₃ H ₄₆ O ₂ = 354.61 [678969-48-7] ([36332-95-3])	≥99% (HPLC) 0.5mg 36,000
	NS480301	2-10 °C	22-Methyltetracosanoic acid (natural) C ₂₅ H ₅₀ O ₂ = 382.66 [678969-50-1] ([36378-43-5])	≥99% (HPLC) 0.5mg 36,000
	NS480401	2-10 °C	24-Methylhexacosanoic acid (natural) C ₂₇ H ₅₄ O ₂ = 410.72 [678969-52-3] ([89838-00-6])	≥99% (HPLC) 0.5mg 36,000
ω-ヒドロキシ 脂肪酸	NS490102 NS490103	-20 °C	30-Hydroxytriacontanoic acid C ₃₀ H ₆₀ O ₃ = 468.80 [52900-18-2]	≥98% (HPLC) 25mg 30,000 100mg 96,000
	NS490202 NS490203	2-10 °C	Methyl 30-hydroxytriacontanoate C ₃₁ H ₆₂ O ₃ = 482.82 [79162-70-2]	≥98% (HPLC) 25mg 15,000 100mg 48,000
	ピロリジジン アルカロイド	NS500100 NS500101	-20 °C	Petasitenine (Fukinotoxin) C ₁₉ H ₂₇ NO ₇ = 381.42 [60102-37-6]
NS500200 NS500201		-20 °C	Neopetasitenine C ₂₁ H ₂₉ NO ₈ = 423.46 [60409-51-0]	≥98% (HPLC) 100µg 12,000 1mg 60,000
NS500300 NS500301		-20 °C	Senkirkine C ₁₉ H ₂₇ NO ₆ = 365.42 [2318-18-5]	≥98% (HPLC) 100µg 12,000 1mg 100,000
カラマツ Larch		NS510102	-20 °C	(+)- <i>trans</i> -Taxifolin C ₁₅ H ₁₂ O ₇ = 304.25 ※吸湿性があり、約10%の水を含む。 [480-18-2]
	NS510201	-20 °C	(+)- <i>trans</i> -Aromadendrin C ₁₅ H ₁₂ O ₆ = 288.25 ※吸湿性があり、約10%の水を含む。 [480-20-6]	≥99.5% (HPLC), ≥99% ee (HPLC) 1mg 12,000
シークワサー Citrus depressa	NS520102 NS520103	-20 °C	Nobiletin C ₂₁ H ₂₂ O ₈ = 402.40 [478-01-3]	≥99% (HPLC) 10mg 20,000 100mg 120,000
	NS520202 NS520203	-20 °C	Tangeretin C ₂₀ H ₂₀ O ₇ = 372.37 [481-53-8]	≥99% (HPLC) 10mg 20,000 100mg 120,000
	NS520302	-20 °C	Sinensetin C ₂₀ H ₂₀ O ₇ = 372.37 [2306-27-6]	≥99% (HPLC) 10mg 36,000
	パセリ Parsley	NS530101 NS530102	-20 °C	Apiin C ₂₆ H ₂₈ O ₁₄ = 564.49 [26544-34-3]
NS540102 NS540103		-20 °C	Myricitrin C ₂₁ H ₂₀ O ₁₂ = 464.38 [17912-87-7]	≥99% (HPLC) 10mg 18,000 100mg 120,000

Source et. al.	Code No.	Storage Conditions	Product Name · Purity · Molecular Formula = Molecular Weight · [CAS Registry Number]	Quantity · Price (JPY)
ニンジン	NS550101	-20°C	Ginsenoside Ra ₁ C ₅₈ H ₉₈ O ₂₅ = 1211.38 [83459-41-0] ≥99%(HPLC)	1mg 36,000
	NS550201	-20°C	Ginsenoside Ra ₂ C ₅₈ H ₉₈ O ₂₅ = 1211.38 [83459-42-1] ≥99%(HPLC)	1mg 36,000
	NS550301	-20°C	Ginsenoside Ra ₃ C ₅₀ H ₁₀₀ O ₂₇ = 1241.41 [90985-77-6] ≥99%(HPLC)	1mg 36,000
	NS550402	-20°C	Ginsenoside Rb ₁ C ₅₄ H ₉₂ O ₂₃ = 1109.29 [41753-43-9] ≥99%(HPLC)	10mg 18,000
	NS550502	-20°C	Ginsenoside Rb ₂ C ₅₃ H ₉₀ O ₂₂ = 1079.27 [11021-13-9] ≥99%(HPLC)	5mg 16,000
	NS550601	-20°C	Ginsenoside Rb ₃ C ₅₃ H ₉₀ O ₂₂ = 1079.27 [68406-26-8] ≥99%(HPLC)	1mg 16,000
	NS550702	-20°C	Ginsenoside Rc C ₅₃ H ₉₀ O ₂₂ = 1079.27 [11021-14-0] ≥99%(HPLC)	5mg 20,000
	NS550801	-20°C	Ginsenoside Rd C ₄₈ H ₈₂ O ₁₈ = 947.15 [52705-93-8] ≥98%(HPLC)	1mg 20,000
	NS550901	-20°C	Malonyl-ginsenoside Rb ₁ C ₅₇ H ₉₄ O ₂₅ = 1195.34 [88140-34-5] ≥98%(HPLC)	1mg 36,000
	NS551001	-20°C	Malonyl-ginsenoside Rb ₂ C ₅₆ H ₉₂ O ₂₅ = 1165.31 [88156-44-9] ≥98%(HPLC)	1mg 36,000
	NS551101	-20°C	Malonyl-ginsenoside Rc C ₅₆ H ₉₂ O ₂₅ = 1165.31 [88140-36-7] ≥98%(HPLC)	1mg 36,000
	NS552102	-20°C	Ginsenoside Re C ₄₈ H ₈₂ O ₁₈ = 947.15 [52286-59-6] ≥99%(HPLC)	5mg 20,000
	NS552202	-20°C	Ginsenoside Rf C ₄₂ H ₇₂ O ₁₄ = 801.01 [52286-58-5] ≥99%(HPLC)	5mg 18,000
	NS552301	-20°C	20-Glucoginsenoside Rf C ₄₈ H ₈₂ O ₁₉ = 963.15 [68406-27-9] ≥99%(HPLC)	1mg 36,000
	NS552402	-20°C	Ginsenoside Rg ₁ C ₄₂ H ₇₂ O ₁₄ = 801.01 [22427-39-0] ≥99%(HPLC)	10mg 24,000
	NS552501	-20°C	Notoginsenoside R ₁ C ₄₇ H ₈₄ O ₁₈ = 933.13 [80418-24-2] ≥99%(HPLC)	1mg 18,000
	NS552601	-20°C	Notoginsenoside R ₂ C ₄₁ H ₇₀ O ₁₉ = 770.99 [80418-25-3] ≥99%(HPLC)	1mg 24,000
NS553102	-20°C	Ginsenoside Ro C ₄₈ H ₇₆ O ₁₉ = 957.11 [34367-04-9] ≥99%(HPLC)	5mg 32,000	
カロテノイド	NS560101 NS560102	-20 °C	β-Cryptoxanthin C ₄₀ H ₅₆ O = 552.87 [472-70-8] ≥99%(HPLC)	1mg 30,000 10mg 150,000

Source et. al.	Code No.	Storage Conditions	Product Name · Purity · Molecular Formula = Molecular Weight · [CAS Registry Number]	Quantity · Price (JPY)
ステロール、 酸化ステロール	NS570102	-20°C	7-Ketocholesterol C ₂₇ H ₄₄ O ₂ = 400.64 [566-28-9]	≥99%(HPLC) 10mg 16,000
	NS570202	-20°C	7α-Hydroxycholesterol C ₂₇ H ₄₆ O ₂ = 402.65 [566-26-7]	≥99%(HPLC) 5mg 32,000
	NS570302	-20°C	7β-Hydroxycholesterol C ₂₇ H ₄₆ O ₂ = 402.65 [566-27-8]	≥99%(HPLC) 5mg 32,000
	NS570401	-20°C	7α-Hydroperoxycholesterol C ₂₇ H ₄₆ O ₃ = 418.65 [2846-29-9]	≥99%(HPLC) 1mg 36,000
	NS570501	-20°C	7β-Hydroperoxycholesterol C ₂₇ H ₄₆ O ₃ = 418.65 [36871-91-7]	≥99%(HPLC) 1mg 36,000
	NS571102	-20°C	β-Sitosterol C ₂₉ H ₅₀ O = 414.71 [83-46-5]	≥99%(HPLC) 5mg 20,000
	NS571202	-20°C	β-Sitosterol acetate C ₃₁ H ₅₂ O ₂ = 456.74 [915-05-9]	≥99%(HPLC) 5mg 20,000
	NS571302	-20°C	7-Keto-β-sitosterol C ₂₉ H ₄₈ O ₂ = 428.69 [2034-74-4]	≥99%(HPLC) 5mg 36,000
	NS571401	-20°C	7α-Hydroxy-β-sitosterol C ₂₉ H ₅₀ O ₂ = 430.71 [34427-61-7]	≥99%(HPLC) 1mg 36,000
	NS571501	-20°C	7β-Hydroxy-β-sitosterol C ₂₉ H ₅₀ O ₂ = 430.71 [15140-59-7]	≥99%(HPLC) 1mg 36,000
	NS572102	-20°C	Stigmasterol C ₂₉ H ₄₈ O = 412.69 [83-48-7]	≥99%(HPLC) 5mg 18,000
	NS572202	-20°C	Stigmasterol acetate C ₃₁ H ₅₀ O ₂ = 454.73 [4651-48-3]	≥99%(HPLC) 5mg 18,000
	NS572302	-20°C	7-Ketostigmasterol C ₂₉ H ₄₆ O ₂ = 426.67 [36449-99-7]	≥99%(HPLC) 5mg 36,000
	NS572401	-20°C	7α-Hydroxystigmasterol C ₂₉ H ₄₈ O ₂ = 428.69 [64998-19-2]	≥99%(HPLC) 1mg 36,000
	NS572501	-20°C	7β-Hydroxystigmasterol C ₂₉ H ₄₈ O ₂ = 428.69 [64998-20-5]	≥99%(HPLC) 1mg 36,000
	NS573101	-20°C	Campesterol C ₂₈ H ₄₈ O = 400.68 [474-62-4]	≥99%(HPLC) 1mg 40,000
	NS573201	-20°C	Campesterol acetate C ₃₀ H ₅₀ O ₂ = 442.72 [1900-53-4]	≥99%(HPLC) 1mg 40,000
	NS573301	-20°C	7-Ketocampesterol C ₂₈ H ₄₆ O ₂ = 414.66 [55396-22-0]	≥99%(HPLC) 1mg 60,000
	NS573401	-20°C	7α-Hydroxycampesterol C ₂₈ H ₄₈ O ₂ = 416.68 [64780-73-0]	≥99%(HPLC) 1mg 70,000
	NS573501	-20°C	7β-Hydroxycampesterol C ₂₈ H ₄₈ O ₂ = 416.68 [64780-91-2]	≥99%(HPLC) 1mg 70,000
NS574101	-20°C	Dihydrobrassicasterol C ₂₈ H ₄₈ O = 400.68 [4651-51-8]	≥99%(HPLC) 1mg 50,000	
NS574201	-20°C	Dihydrobrassicasterol acetate C ₃₀ H ₅₀ O ₂ = 442.72 [4651-52-9]	≥99%(HPLC) 1mg 50,000	
NS574301	-20°C	7-Ketodihydrobrassicasterol C ₂₈ H ₄₆ O ₂ = 414.66 [156767-69-0]	≥99%(HPLC) 1mg 50,000	
NS574401	-20°C	7α-Hydroxydihydrobrassicasterol C ₂₈ H ₄₈ O ₂ = 416.68 [71486-05-0]	≥99%(HPLC) 1mg 80,000	
NS574501	-20°C	7β-Hydroxydihydrobrassicasterol C ₂₈ H ₄₈ O ₂ = 416.68 [71486-04-9]	≥99%(HPLC) 1mg 80,000	
たもぎ茸	NS580102 NS580103 <i>new</i>	2-10 °C	L-(+)-Ergothioneine C ₉ H ₁₅ N ₃ O ₂ S = 229.30 [497-30-3]	≥98% (HPLC) 10mg 16,000 100mg 120,000
シゴカノ エゾウコギ	NS600102 <i>new</i>	2-10 °C	Isofraxidin C ₁₁ H ₁₀ O ₅ = 222.19 [486-21-5]	≥99% (HPLC) 10mg 16,000
	<i>new</i>		Eleutheroside B C ₁₇ H ₂₄ O ₉ = 372.37 [118-34-3]	≥99% (HPLC)
	<i>new</i>		Eleutheroside B ₁ C ₁₇ H ₂₀ O ₁₀ = 384.34 [16845-16-2]	≥99% (HPLC)
ハッカ	NS610102 <i>new</i>	-20°C	ジヒドロナフトレン誘導体 [Salvianolic acid L] C ₃₀ H ₃₀ O ₁₆ = 718.62 [147568-62-5, 389065-74-1]	≥99% (TLC) 10mg 15,000
シベット	NS620102 <i>new</i>	2-10 °C	Civetone C ₁₇ H ₃₀ O = 250.42 [542-46-1]	≥99% (TLC) 10mg 36,000
南天実	NS640102 <i>new</i>	2-10 °C	O-Methyl domesticine [Nantenine] C ₂₀ H ₂₁ NO ₄ = 339.39 [2565-01-7, 42971-15-3]	≥99% (HPLC) 10mg 18,000

Source et. al.	Code No.	Storage Conditions	Product Name · Purity · Molecular Formula=Molecular Weight · [CAS Registry Number]	Quantity · Price (JPY)
プベルル酸	NS650101 NS650102 <i>new</i>	2-10 °C	Puberulic acid ≥98% (HPLC) C ₈ H ₆ O ₆ =198.13 [99-23-0]	2mg 25,000 10mg 50,000
アゼライン酸	NS660104 <i>new</i>	2-10 °C	Azelaic acid ≥99% (HPLC) C ₉ H ₁₆ O ₄ =188.22 [123-99-9]	1g 9,000

Source et. al.	Code No.	Storage Conditions	Product Name · Purity · Molecular Formula = Molecular Weight · [CAS Registry Number]	Quantity · Price (JPY)
シアル酸誘導体等	SA001002 SA001010 SA001025	-20 °C	<i>N</i> Acetylneuraminic acid (NANA ; Sialic acid) ≥99% (HPLC) crystal C ₁₁ H ₁₉ NO ₉ = 309.27 [131-48-6]	2g 16,000 10g 26,000 25g 60,000
	SA002002 SA002010	-20 °C	<i>N</i> Acetylneuraminic acid, methyl ester (Sialic acid, methyl ester) ≥95% (TLC) C ₁₂ H ₂₁ NO ₉ = 323.30 [22900-11-4]	2g 24,000 10g 96,000
	SA003002 SA003010	-20 °C	<i>N</i> Acetylneuraminic acid, 2,4,7,8,9-pentaacetate (Sialic acid, 2,4,7,8,9-pentaacetate) ≥95% (TLC) C ₂₁ H ₂₉ NO ₁₄ = 519.45 [4887-11-0]	2g 24,000 10g 96,000
	SA004001 SA004010	-20 °C	<i>N</i> Acetylneuraminic acid, methyl ester, 2,4,7,8,9-pentaacetate (Sialic acid, methyl ester, 2,4,7,8,9-pentaacetate) ≥95% (TLC) C ₂₂ H ₃₁ NO ₁₄ = 533.48 [73208-82-9]	1g 30,000 10g 210,000
	SA005001 SA005010	冷暗所	<i>N</i> Acetyl-2-phenylthionneuraminic acid, methyl ester, 4,7,8,9-tetraacetate (2-(Sph)-sialic acid, methyl ester, 4,7,8,9-tetraacetate) (α, β mix) ≥98% (TLC) C ₂₆ H ₃₃ NO ₁₂ S = 583.61 [155155-64-9]	1g 40,000 10g 320,000
	SA006001 SA006005	2-10 °C	<i>N</i> Iodosuccinimide ¹⁾²⁾ ≥99% (TLC) C ₄ H ₄ INO ₂ = 224.98 [516-12-1]	1g 18,000 5g 36,000
	SA007010 SA007050 SA007100	-20 °C	2- <i>O</i> -Methyl α- <i>D</i> - <i>N</i> acetylneuraminic acid ≥98% (TLC) C ₁₂ H ₂₁ NO ₉ = 323.30 [50930-22-8]	10mg 22,000 50mg 88,000 100mg 154,000
	SA008010 SA008050 SA008100	-20 °C	2- <i>O</i> -Phenyl α- <i>D</i> - <i>N</i> acetylneuraminic acid ≥98% (TLC) C ₁₇ H ₂₃ NO ₉ = 385.37 [15964-32-6]	10mg 30,000 50mg 120,000 100mg 210,000
	SA010100	2-10 °C	Methyl 2,3-didehydro-4,7,8,9-tetra- <i>O</i> -acetyl- <i>N</i> acetylneuraminic acid ≥98% (TLC) C ₂₀ H ₂₇ NO ₁₂ = 473.42 [73960-72-2]	100mg 30,000
	SA011001	冷暗所	<i>N</i> Acetyl-2-phenylthio-α-neuraminic acid, methyl ester, 4,7,8,9-tetraacetate (2-(Sph)-α-sialic acid, methyl ester, 4,7,8,9-tetraacetate) ≥98% (TLC) C ₂₆ H ₃₃ NO ₁₂ S = 583.61 [118977-26-7]	1g 60,000
グルコース誘導体	GL001010	2-10 °C	4-Methoxyphenyl 2,3,4,6-tetra- <i>O</i> -acetyl-β- <i>D</i> -glucopyranoside ≥98% (TLC) C ₂₁ H ₂₅ O ₁₁ = 454.42 [14581-81-8]	10g 15,000
	GL002005	2-10 °C	4-Methoxyphenyl β- <i>D</i> -glucopyranoside ≥98% (TLC) C ₁₃ H ₁₈ O ₇ = 286.28 [6032-32-2]	5g 22,000
	GL003005	2-10 °C	Phenyl 2,3,4,6-tetra- <i>O</i> -acetyl-1-thio-β- <i>D</i> -glucopyranoside ≥98% (TLC) C ₂₀ H ₂₁ O ₉ S = 440.46 [23661-28-1]	5g 22,000
	GL004001 GL004005	2-10 °C	Phenyl 1-thio-β- <i>D</i> -glucopyranoside ≥98% (TLC) C ₁₂ H ₁₆ O ₅ S = 272.32 [2936-70-1]	1g 25,000 5g 100,000
	GL005001 GL005005	2-10 °C	Ethyl 2,3,4,6-tetra- <i>O</i> -acetyl-1-thio-β- <i>D</i> -glucopyranoside ≥98% (TLC) C ₁₆ H ₂₁ O ₉ S = 392.42 [52645-73-5]	1g 15,000 5g 60,000
	GL006001 GL006005	2-10 °C	Ethyl 1-thio-β- <i>D</i> -glucopyranoside ≥98% (TLC) C ₈ H ₁₆ O ₅ S = 224.28 [7473-36-1]	1g 30,000 5g 120,000
	ガラクトース誘導体	GA001010	2-10 °C	4-Methoxyphenyl 2,3,4,6-tetra- <i>O</i> -acetyl-β- <i>D</i> -galactopyranoside ≥98% (TLC) C ₂₁ H ₂₅ O ₁₁ = 454.42 [2872-65-3]
GA002005		2-10 °C	4-Methoxyphenyl β- <i>D</i> -galactopyranoside ≥98% (TLC) C ₁₃ H ₁₈ O ₇ = 286.28 [3150-20-7]	5g 24,000
GA003002		2-10 °C	4-Methoxyphenyl 2,6-di- <i>O</i> -benzyl-β- <i>D</i> -galactopyranoside ≥98% (TLC) C ₂₇ H ₃₀ O ₇ = 466.52 [159922-50-6]	2g 24,000
GA004001		2-10 °C	4-Methoxyphenyl 2,3-di- <i>O</i> -benzyl-β- <i>D</i> -galactopyranoside ≥98% (TLC) C ₂₇ H ₃₀ O ₇ = 466.52	1g 20,000
GA005005		2-10 °C	Phenyl 2,3,4,6-tetra- <i>O</i> -acetyl-1-thio-β- <i>D</i> -galactopyranoside ≥98% (TLC) C ₂₀ H ₂₁ O ₉ S = 440.47 [24404-53-3]	5g 28,000
GA006001 GA006005		2-10 °C	Phenyl 1-thio-β- <i>D</i> -galactopyranoside ≥98% (TLC) C ₁₂ H ₁₆ O ₅ S = 272.32 [16758-34-2]	1g 30,000 5g 120,000
GA007001		2-10 °C	2-(Trimethylsilyl)ethyl β- <i>D</i> -galactopyranoside ≥98% (TLC) C ₁₁ H ₂₄ O ₆ Si = 280.39 [117252-95-6]	1g 24,000
ラクトース誘導体	LA001500 LA001001	2-10 °C	4-Methoxyphenyl 4- <i>O</i> -β- <i>D</i> -galactopyranosyl-β- <i>D</i> -glucopyranoside ≥98% (TLC) C ₁₉ H ₂₈ O ₁₂ = 448.42 [150412-80-9]	500mg 20,000 1g 32,000
	LA002500 LA002001	2-10 °C	4-Methoxyphenyl 2,2',3,6,6'-penta- <i>O</i> -benzyl-4- <i>O</i> -β- <i>D</i> -galactopyranosyl-β- <i>D</i> -glucopyranoside ≥98% (TLC) C ₅₄ H ₅₈ O ₁₂ = 899.03 [358681-61-5]	500mg 36,000 1g 60,000
	LA003500 LA003001	2-10 °C	4-Methoxyphenyl 2,2',3,3',6,6'-penta- <i>O</i> -benzyl-4- <i>O</i> -β- <i>D</i> -galactopyranosyl-β- <i>D</i> -glucopyranoside ≥98% (TLC) C ₅₄ H ₅₈ O ₁₂ = 899.03	500mg 36,000 1g 60,000
	LA004500	2-10 °C	2-(Trimethylsilyl)ethyl 4- <i>O</i> -β- <i>D</i> -galactopyranosyl-β- <i>D</i> -glucopyranoside ≥98% (TLC) C ₁₇ H ₃₄ O ₁₁ Si = 442.53 [115969-51-2]	500mg 24,000
	LA005500	2-10 °C	Benzyl 4- <i>O</i> -β- <i>D</i> -galactopyranosyl-β- <i>D</i> -glucopyranoside ≥98% (TLC) C ₁₉ H ₂₈ O ₁₁ = 432.42 [18404-72-3]	500mg 24,000

Source et. al.	Code No.	Storage Conditions	Product Name · Purity · Molecular Formula=Molecular Weight · [CAS Registry Number]	Quantity · Price (JPY)
コロミン酸塩	SA009100 SA009001	2-10 °C	Colominic acid, sodium salt from <i>E. coli</i> ≥98% [70431-34-4]	100mg 30,000 1g 100,000
ヒアルロン酸	HA001005 HA001025	2-10 °C	Hyaluronic Acid from <i>Streptococcus</i> for Biochem. [9004-61-9]	5g 15,000 25g 60,000
	HA002005 HA002025	2-10 °C	Hyaluronic Acid Sodium Salt from <i>Streptococcus</i> for Biochem. [9067-32-7]	5g 15,000 25g 60,000
ジシアル酸等	SA012100	-20 °C	<i>N</i> -Acetylneuraminic acid, dimer ($\alpha, 2 \rightarrow 8$) ≥95% (TLC) (<i>N</i> -acetyl-8- <i>O</i> -(<i>N</i> -acetyl- α -neuraminosyl)-neuraminic acid, disodium salt) C ₂₂ H ₃₄ N ₂ Na ₂ O ₁₇ =644.49 [149331-75-9]	100mg 25,000
	SA013100	-20 °C	<i>N</i> -Acetyl-8- <i>O</i> -(<i>N</i> -acetyl-4',7,8,9'-tetra- <i>O</i> -acetyl- α -neuraminosyl)-2- <i>S</i> -phenyl-2-thio-neuraminic acid, 1',9-ester, 1-methyl-ester, 4,7-diacetate C ₄₁ H ₅₂ N ₂ O ₂₁ S=940.92 [158111-03-6]	100mg 38,000
シアル酸含有糖鎖	SG001010 SG001100 SG001001	2-10 °C	Protected 6'- α -Sialylgalactose; Protected NeuAc α (2→6)-D-Gal ≥98% (TLC) C ₄₇ H ₅₇ NO ₁₉ =939.95	10mg 20,000 100mg 90,000 1g 630,000
	SG002010 SG002100 SG002001	2-10 °C	Protected 3'- α -Sialylgalactose; Protected NeuAc α (2→3)-D-Gal ≥98% (TLC) C ₄₇ H ₅₇ NO ₁₉ =939.95	10mg 20,000 100mg 90,000 1g 630,000
	SG003010 SG003100	2-10 °C	Protected 6'- β -Sialylgalactose; Protected NeuAc β (2→6)-D-Gal ≥98% (TLC) C ₄₇ H ₅₇ NO ₁₉ =939.95	10mg 30,000 100mg 180,000
	SL001010 SL001100 SL001001	2-10 °C	Protected 6'- α -Sialyllactose; Protected NeuAc α (2→6)-D-Gal- β (1→4)-D-Glc ≥98% (TLC) C ₇₄ H ₈₅ NO ₂₄ =1372.46	10mg 28,000 100mg 140,000 1g 980,000
	SL002010 SL002100 SL002001	2-10 °C	Protected 3'- α -Sialyllactose; Protected NeuAc α (2→3)-D-Gal- β (1→4)-D-Glc ≥98% (TLC) C ₇₄ H ₈₅ NO ₂₄ =1372.46	10mg 28,000 100mg 140,000 1g 980,000
	SL003010 SL003100	2-10 °C	Protected 6'- β -Sialyllactose; Protected NeuAc β (2→6)-D-Gal- β (1→4)-D-Glc ≥98%(TLC) C ₇₄ H ₈₅ NO ₂₄ =1372.46	10mg 45,000 100mg 270,000
	SL006010 SL006050	-20 °C	3'-Sialyllactose sodium salt ≥99%(TLC) C ₂₃ H ₃₈ NO ₁₉ Na=655.53 [128596-80-5, Component: 35890-38-1]	10mg 10,000 50mg 20,000
	SL007010 SL007050	-20 °C	6'-Sialyllactose sodium salt ≥99%(TLC) C ₂₃ H ₃₈ NO ₁₉ Na=655.53 [157574-76-0, Component: 35890-39-2]	10mg 10,000 50mg 20,000
	SL008005 SL008010	-20 °C	Disialyllactose sodium salt ≥99%(TLC) C ₃₄ H ₅₄ N ₂ O ₂₇ Na ₂ =968.77 []	5mg 30,000 10mg 54,000

分離用シリカゲル系担体

1. 高純度 (99.99%) 球形シリカゲル Namsil NX-300H

高純度シリカゲルはフェノール性天然物の分離に威力を発揮することから、近年 HPLC 充填カラムに広く用いられるようになった。当社では回収率よく高度に分級（粒揃え）することにより安価なオープンカラム用の高純度 (99.99%) シリカゲルの製造法を確立した。高純度シリカゲルは金属イオンを殆ど含まないため、1) シャープなピークを与える。2) 目的物の殆どを回収できる。3) 原点に着色物質などが残らず、数十回以上再利用できる。

品名	価格 (円)
Namsil NX-300H (300mesh)	100g 27,000 1kg 180,000

2. 破砕形シリカゲル Namgel NA-200H, 300H

200mesh, 300mesh の一般用の特性シリカゲルで、高流速を特徴としている。

品名	価格 (円)
Namgel NA-200H (200mesh)	1kg 8,000
Namgel NA-300H (300mesh)	1kg 10,000

3. 極性化合物用 破砕形シリカゲル Namgel MAN-200H, 300H

極性化合物の分離を目的として特別に開発したゲルで、メタノールなどを含む極性溶媒で分離する場合に特に威力を示す。

品名	価格 (円)
Namgel NAM-200H (200mesh)	1kg 12,000
Namgel NAM-300H (300mesh)	1kg 16,000

*各種サイズのシリカゲル、蛍光剤入り、蛍光検出用カラムの在庫がございますので弊社までお問い合わせ下さい。

瞬速分析・分取 2 μ HPLC 充填カラム

〒501-1121 岐阜市古市場 840

長良サイエンス (株)

社長 中塚進一

Tel:058-234-4257 Fax: 058-234-4724

E-mail:nagara@nsgifu.jp URL:http://www.nsgifu.jp

HPLC等のクロマトグラフィーに用いる充填剤は、その粒度分布により使用圧力や分離性能が大きく影響する。弊社では独自の精密分級(粒前え)技術を開発して、目詰まりの原因となる微粒子と分離能低下の原因となる粗大粒子をカットすることに成功しました。これにより高分離能で使用圧力が20%以上低く、長寿命の充填カラムを開発できた。その結果、~1000気圧の超高压が必須とされてきた2 μ ODSシリカゲルの充填カラムが、通常のHPLC装置(250気圧以下)、数分で分析・分取(~数十mg)が可能となりました。

①JASIS (分析展、科学機器展、幕張メッセ 2012.9)


②食品開発展 (東京ビッグサイト、2012.10)

③日本農芸化学会年会 (東北大学、2013.3)

「瞬速2 μ ODSシリカゲルのHPLC充填カラムの開発」

2 μ ODS (4.6mmID×50mm)

舞茸由来グルコシルセラミド混合物標準品



充填カラムの種類	コード番号	サイズ	使用目的	装置	標準流量	分析時間	価格 (円)
Nagara ODS-2 炭素含量 17%	NS ODS2-020050	2.0mm×50mm	微量分析	HPLC*	0.2ml/min	3min	40,000
	NS ODS2-030050	3.0mm×50mm	分析	HPLC*	0.4ml/min	3min	43,000
	NS ODS2-046050	4.6mm×50mm	分析/分取	HPLC	1ml/min	3min	一時中止、再開未定
	NS ODS2-100050	10mm×50mm	分取	HPLC	5ml/min	3min	一時中止、再開未定

*注 これらの充填カラムでは、内容量が極めて小さいために装置配管中での拡散を無視できません。汎用のHPLC装置で3mm ϕ ×50mmカラムを使用すると化合物の分析ピークがブロードになり、2mm ϕ ×50mmでは相当崩れます。これを改善するため、弊社では独自に改造した微量用HPLC装置やUHPLC装置を使用しています。